

# FINITE ABUNDANT SEMIGROUPS IN WHICH THE IDEMPOTENTS FORM A SUBSEMIGROUP\*

John Fountain  
Department of Mathematics, University of York,  
Heslington, York YO10 5DD, U.K.  
**e-mail** : jbf1@york.ac.uk

and

Gracinda M. S. Gomes  
Centro de Álgebra da Universidade de Lisboa  
Avenida Prof Gama Pinto 2  
1649-003 Lisboa, Portugal  
and  
Departamento de Matemática  
Faculdade de Ciências  
Universidade de Lisboa  
1746-016 Lisboa, Portugal  
**e-mail** : ggomes@cii.fc.ul.pt

**ABSTRACT.** We consider certain abundant semigroups in which the idempotents form a subsemigroup, and which we call bountiful semigroups. We find a simple criterion for a finite bountiful semigroup to be a member of the join of the pseudovarieties of finite groups and finite aperiodic semigroups.

## INTRODUCTION

In the 1970s Schützenberger posed the problem of finding a characterisation of the semigroups in the pseudovariety  $\mathbf{A} \vee \mathbf{G}$  where  $\mathbf{A}$  is the pseudovariety of all finite aperiodic semigroups, and  $\mathbf{G}$  is the pseudovariety of all finite groups. McAlister answered this question for orthodox semigroups [13] by showing that a finite orthodox semigroup  $S$  is in  $\mathbf{A} \vee \mathbf{G}$  if and only if  $\mathcal{H}$  is a congruence on  $S$ . He subsequently generalised this result to obtain a characterisation of the regular semigroups in  $\mathbf{A} \vee \mathbf{G}$  [14]. Recently, Steinberg [21] has shown that McAlister's results can be deduced from work of Rhodes and Tilson [18] and that this approach allows McAlister's theorems to be extended to a wider class of

---

2000 *Mathematics Subject Classification.* Primary: 20M10.

*Key words and phrases.* covers of semigroups, pseudovarieties of semigroups.

\*This work was partially supported by "Financiamento Programático of CAUL" and project POCTI/32440/MAT/2000 of FCT and FEDER.

semigroups. Our purpose in this paper is to extend the result for orthodox semigroups in a different direction.

Recall that the relation  $\mathcal{R}^*$  on a semigroup  $S$  is defined by  $a\mathcal{R}^*b$  if and only if  $a$  and  $b$  are  $\mathcal{R}$ -related in an extension of  $S$ . The dual of  $\mathcal{R}^*$  is  $\mathcal{L}^*$ , and  $\mathcal{H}^* = \mathcal{R}^* \cap \mathcal{L}^*$ . A semigroup is *abundant* if every  $\mathcal{R}^*$ -class and every  $\mathcal{L}^*$ -class contains an idempotent. We say that an abundant semigroup is *bountiful* if it is idempotent-connected (see Section 1 for the definition) and its idempotents form a subsemigroup. Let  $\mu$  be the greatest congruence contained in  $\mathcal{H}^*$ . We now give the main result of the paper.

**Theorem 1.** *Let  $S$  be a finite bountiful semigroup. Then  $S \in \mathbf{A} \vee \mathbf{G}$  if and only if  $\mathcal{H} \subseteq \mu$ .*

In the following section, we recall some basic results related to abundant semigroups. Following [1], we say that an idempotent-connected abundant semigroup in which the idempotents generate a regular subsemigroup is *concordant*. For a concordant semigroup  $S$  with set of idempotents  $E(S)$ , El Qallali and Fountain [2] gave a representation of  $S$  as a full subsemigroup of the Hall-Nambooripad semigroup  $T_{\langle E(S) \rangle}$  with kernel  $\mu$ . We use this to prove the main result of Section 2: if  $S$  is a concordant semigroup in  $\mathbf{A} \vee \mathbf{G}$ , then  $\mathcal{H} \subseteq \mu$  on  $S$ . Our proof also requires the corresponding result for regular semigroups, due to McAlister [13]. In the final section, we complete the proof of the main theorem by showing that if  $S$  is a bountiful semigroup satisfying  $\mathcal{H} \subseteq \mu$ , then  $S \in \mathbf{A} \vee \mathbf{G}$ . Our proof is modelled on that for the orthodox case; we require a preliminary result to the effect that a finite bountiful semigroup has a finite bountiful  $E$ -unitary cover of a special type. The existence of bountiful  $E$ -unitary covers for bountiful semigroups has recently been established by Simmons [20], but his proof does not give finite covers for finite semigroups. Our proof mimics that of McAlister [13] for the orthodox case, and uses the fact that a finite ample semigroup has a finite proper cover, a result due to the authors [6]. That ample semigroups have proper covers was first shown by Lawson [11], and is also a consequence of Simmons' work on bountiful semigroups. However, their proofs do not give finite covers for finite semigroups.

## 1. PRELIMINARIES

For basic semigroup notation and terminology, we follow [9]. In particular,  $E(S)$  denotes the set of idempotents of a semigroup  $S$ . We recall the following alternative characterisation of  $\mathcal{R}^*$  from [12] and [16] which we shall use without further mention.

**Lemma 1.1.** *The following are equivalent for elements  $a, b$  of a semigroup  $S$ :*

- (1)  $a\mathcal{R}^*b$ ,
- (2) for all  $x, y \in S^1$ ,  $xa = ya$  if and only if  $xb = yb$ .

This condition is simplified when one of the elements involved is an idempotent.

**Corollary 1.2.** *Let  $a$  be an element of a semigroup  $S$ , and  $e \in E(S)$ . Then the following are equivalent:*

- (1)  $a\mathcal{R}^*e$ ,
- (2)  $ea = a$  and for all  $x, y \in S^1$ ,  $xa = ya$  implies  $xe = ye$ .

We remark that  $\mathcal{R}^*$  is a left congruence, and  $\mathcal{L}^*$  is a right congruence. Also on any semigroup  $S$  we have  $\mathcal{R} \subseteq \mathcal{R}^*$ . It is well known and easy to see that if  $a, b \in S$  are regular, then  $a\mathcal{R}^*b$  if and only if  $a\mathcal{R}b$ . In particular, if  $S$  is regular, then  $\mathcal{R}^* = \mathcal{R}$ . If there is any danger of ambiguity, we use  $\mathcal{R}^*(S)$ , etc. to denote the relation  $\mathcal{R}^*$  on  $S$ . The  $\mathcal{R}^*$ -class of  $a \in S$  will be denoted by  $R_a^*$  or  $R_a^*(S)$ , and corresponding notation is used for  $\mathcal{L}^*$ - and  $\mathcal{H}^*$ -classes.

A semigroup in which each  $\mathcal{R}^*$ -class and each  $\mathcal{L}^*$ -class contains an idempotent is said to be *abundant*. From [2] we have the following lemma.

**Lemma 1.3.** *Let  $U$  be an abundant subsemigroup of an abundant semigroup  $S$  such that the idempotents of  $U$  form an order-ideal of those of  $S$ . Then*

$$\mathcal{R}^*(U) = \mathcal{R}^*(S) \cap (U \times U).$$

The next corollary is an easy consequence of the lemma and its dual.

**Corollary 1.4.** *Let  $S$  be an abundant semigroup. Then every full subsemigroup of  $S$  is abundant, and if  $e \in E(S)$ , then  $eSe$  is abundant.*

We say that a homomorphism  $\varphi : S \rightarrow T$  of abundant semigroups is *good* if, for all elements  $a, b$  of  $S$ , we have  $a\mathcal{R}^*(S)b$  implies  $a\varphi\mathcal{R}^*(T)b\varphi$ , and  $a\mathcal{L}^*(S)b$  implies  $a\varphi\mathcal{L}^*(T)b\varphi$ . A congruence  $\rho$  on an abundant semigroup  $S$  is *good* if the natural homomorphism  $S \rightarrow S/\rho$  is good. We remark that any homomorphism with regular domain is good.

Let  $S$  be an abundant semigroup and  $B$  be the subsemigroup generated by  $E(S)$ . We say that  $S$  is *idempotent-connected* (IC) when for each element  $a$  of  $S$  and some idempotents  $e, f$  in  $R_a^*$  and  $L_a^*$  respectively, there is a bijection  $\alpha : \langle E(eBe) \rangle \rightarrow \langle E(fBf) \rangle$  satisfying  $xa = a(x\alpha)$  for all  $x \in \langle E(eBe) \rangle$ . We remark that in [2] it is shown that the word ‘‘some’’ can be replaced by ‘‘all’’, and that the bijection  $\alpha$  is an isomorphism. It is also worth mentioning that any regular semigroup  $S$  is IC since, for any  $a \in S$  and idempotents  $e, f$  in  $R_a$  and  $L_a$ , there is an inverse  $a'$  with  $aa' = e$ ,  $a'a = f$  and we have an isomorphism  $\alpha : \langle E(aa'Baa') \rangle \rightarrow \langle E(a'aBa'a) \rangle$  with the required property given by  $xa = a'xa$ .

More details about IC abundant semigroups, and alternative formulations of the definition can be found in [1], [2], [10], and [20].

As mentioned in the introduction, the congruence  $\mu$  on an abundant semigroup  $S$  is the largest congruence contained in  $\mathcal{H}^*$ . Hence if  $S$  is regular,  $\mu$  is the largest congruence contained in  $\mathcal{H}$ , and so it is the maximum idempotent separating congruence on  $S$ . Thus our notation is consistent with the standard notation for regular semigroups. When more than one semigroup is involved, we write  $\mu_S$  for the relation  $\mu$  on  $S$ .

## 2. CONCORDANT SEMIGROUPS

A *concordant* semigroup is an IC abundant semigroup in which the idempotents generate a regular semigroup. In this section we prove the following result.

**Proposition 2.1.** *If  $S$  is a finite concordant semigroup in  $\mathbf{A} \vee \mathbf{G}$ , then  $\mathcal{H} \subseteq \mu$ .*

Our approach is to use a representation (due to El Qallali and Fountain [2]) of a concordant semigroup  $S$  in a ‘fundamental’ regular semigroup obtained from  $\langle E(S) \rangle$  by a construction due to Hall [8]. First, we note the following alternative description of concordant semigroups.

**Lemma 2.2.** *An IC abundant semigroup is concordant if and only if the regular elements form a subsemigroup.*

*Proof.* This is immediate by Result 7 of [8]. □

From [1, Lemma 2.4 and Theorem 2.5], we have the following two lemmas, the first being what we might call Lallement’s lemma for concordant semigroups.

**Lemma 2.3.** *Let  $S$  be a concordant semigroup and  $\varphi: S \rightarrow T$  be a good homomorphism. If  $a \in S$  is such that  $a\varphi \in E(T)$ , then  $a\varphi = h\varphi$  for some idempotent  $h \in S$ .*

**Lemma 2.4.** *Let  $S$  be a concordant semigroup. If  $\varphi: S \rightarrow T$  is a surjective good homomorphism, then  $T$  is concordant.*

We make use of results from [8] and [2] for which we need the following semigroup constructed in [8] (see also [15]); given an idempotent generated regular semigroup  $\langle E \rangle$  with set of idempotents  $E$ , Hall constructs a regular semigroup  $T_{\langle E \rangle}$  such that  $\langle E(T_{\langle E \rangle}) \rangle$  is isomorphic to  $\langle E \rangle / \mu_{\langle E \rangle}$ . Moreover,  $T_{\langle E \rangle}$  is *fundamental*, that is, the congruence  $\mu_{T_{\langle E \rangle}}$  is trivial. The following theorem is due to Hall [8] in the regular case, and was extended to concordant semigroups in [2]. The construction of  $T_{\langle E(S) \rangle}$  does not play a part in the arguments of this paper; all we need is that it exists, and some of its properties.

**Theorem 2.5.** *Let  $S$  be a concordant semigroup with set of idempotents  $E$  and let  $\langle E \rangle$  be the subsemigroup generated by  $E$ . Then there is a good homomorphism  $\alpha: S \rightarrow T_{\langle E \rangle}$  such that*

- (1)  $\mu = \alpha\alpha^{-1}$ , and
- (2)  $S\alpha$  is a full subsemigroup of  $T_{\langle E \rangle}$ .

*Moreover, every full subsemigroup of  $T_{\langle E \rangle}$  is fundamental.*

Note that as a consequence of the theorem,  $\mu$  is a good congruence on  $S$ .

By Lemma 2.4,  $S\alpha$  is concordant, and so, by Lemma 2.2,  $\text{Reg}(S\alpha)$  is a subsemigroup of  $S\alpha$  which is obviously full. By the theorem,  $S\alpha$  is a full subsemigroup of  $T_{\langle E \rangle}$ , and so  $\text{Reg}(S\alpha)$  is a full regular subsemigroup of  $T_{\langle E \rangle}$ . Hence  $\text{Reg}(S\alpha)$  is fundamental.

Now, if  $S$  is a member of  $\mathbf{A} \vee \mathbf{G}$ , then so is  $\text{Reg}(S\alpha)$ , and so, by Proposition 1.6 of [13],  $\mathcal{H}$  is a congruence on  $\text{Reg}(S\alpha)$ . Thus,  $\mathcal{H} = \mu$  on  $\text{Reg}(S\alpha)$  so that  $\text{Reg}(S\alpha)$  is  $\mathcal{H}$ -trivial.

Now let  $e \in E(S)$ . Then  $e\alpha \in E(S\alpha)$ . By [5, Lemma 1.12], the  $\mathcal{H}^*$ -class  $H_{e\alpha}^*(S\alpha)$  of  $e\alpha$  in  $S\alpha$  is a cancellative subsemigroup of  $S\alpha$ ; but  $S\alpha$  is finite, so  $H_{e\alpha}^*(S\alpha)$  is a group, and hence coincides with the  $\mathcal{H}$ -class of  $e\alpha$  in  $S\alpha$ . Clearly,  $H_{e\alpha}^*(S\alpha) \subseteq \text{Reg}(S\alpha)$ , and it follows easily that it is also the  $\mathcal{H}^*$ -class of  $e\alpha$  in  $\text{Reg}(S\alpha)$ . But  $\text{Reg}(S\alpha)$  is regular, so  $\mathcal{H}^*$  coincides with  $\mathcal{H}$  on  $\text{Reg}(S\alpha)$ , and since  $\text{Reg}(S\alpha)$  is  $\mathcal{H}$ -trivial, we see that  $H_{e\alpha}^*(S\alpha) = \{e\alpha\}$ . Thus all subgroups of  $S\alpha$  are trivial, that is,  $S\alpha \in \mathbf{A}$ . By [17, Proposition 3.4.2],  $S\alpha$  is  $\mathcal{H}$ -trivial, and since homomorphisms map  $\mathcal{H}$ -related elements to  $\mathcal{H}$ -related elements, it follows that, on  $S$ , we have  $\mathcal{H} \subseteq \alpha\alpha^{-1} = \mu$ . This completes the proof of Proposition 2.1.

### 3. BOUNTIFUL SEMIGROUPS

A *bountiful* semigroup is an IC abundant semigroup in which the idempotents form a subsemigroup. Thus a bountiful semigroup is concordant. In this section, we prove that if  $S$  is a bountiful semigroup with  $\mathcal{H} \subseteq \mu$ , then  $S \in \mathbf{A} \vee \mathbf{G}$ . Our proof relies on a result from [6] which we now explain.

An *ample* semigroup is a bountiful semigroup in which the subsemigroup of idempotents is commutative. In an ample semigroup  $S$ , each  $\mathcal{R}^*$ -class and each  $\mathcal{L}^*$ -class contains a unique idempotent. For  $a \in S$ , we denote the idempotent in  $R_a^*$  by  $a^+$ , and that in  $L_a^*$  by  $a^*$ . Thus we may regard an ample semigroup as a (2,1,1)-algebra with unary operations  $+$  and  $*$ . We note that a semigroup homomorphism  $\theta: S \rightarrow T$  of ample semigroups is good if and only if it is a (2,1,1)-algebra homomorphism.

As is well known, every semigroup  $S$  has a minimum cancellative congruence, and we denote this by  $\sigma$ . For regular or finite semigroups,  $\sigma$  is, of course, the minimum group congruence. An explicit description of  $\sigma$  on a bountiful semigroup is given in the next proposition which is due to Simmons [20, Proposition 6].

**Proposition 3.1.** *If  $S$  is a bountiful semigroup, then the minimum cancellative congruence  $\sigma$  on  $S$  is given by:*

$$a\sigma b \text{ if and only if } ea = bf \text{ for some } e, f \in E(S).$$

An ample semigroup is *proper* if  $\mathcal{R}^* \cap \sigma = \iota = \mathcal{L}^* \cap \sigma$ . It follows from [4], and is not difficult to show, that a proper ample semigroup is  $E$ -unitary, and, of course, an inverse semigroup is  $E$ -unitary if and only if it is proper. However, the semigroup of Example 3 in [4] is easily seen to be ample; it is noted in [4] that it is not proper but is  $E$ -unitary.

A proper ample semigroup  $P$  is a *proper cover* of an ample semigroup  $S$  if there is a surjective good homomorphism  $\alpha: P \rightarrow S$  such that  $\alpha$  maps  $E(P)$  isomorphically onto  $E(S)$ . The homomorphism  $\alpha$  is called a *covering homomorphism*. From [6], we have

**Theorem 3.2.** *Every ample semigroup  $S$  has a proper cover which can be taken to be finite if  $S$  is finite.*

As mentioned in the introduction, this result without the finiteness condition was obtained by Lawson [11] and Simmons [20].

Let  $S$  be a bountiful semigroup. For  $e \in E(S)$ , we denote the  $\mathcal{D}$ -class of  $e$  in  $E(S)$  by  $E(e)$ . The relation  $\delta$  on  $S$  is defined by the rule:

$$a\delta b \text{ if and only if } b = eaf \text{ for some } e \in E(h), f \in E(k) \\ \text{where } h \in E(S) \cap R_a^* \text{ and } k \in E(S) \cap L_a^*.$$

It is shown in [3] that if  $\delta$  is a congruence, then it is the minimum ample good congruence on  $S$ , and that  $\mathcal{H}^* \cap \delta = \iota$ . Ronghua [19] and Guo [7] independently proved that  $\delta$  is a congruence on any bountiful semigroup. Putting these results together, we have the following proposition.

**Proposition 3.3.** *A bountiful semigroup  $S$  is a subdirect product of  $S/\mu$  and  $S/\delta$ .*

An  $E$ -unitary cover of a bountiful semigroup  $S$  is an  $E$ -unitary bountiful semigroup  $T$  together with a surjective good homomorphism  $\psi: T \rightarrow S$  which maps  $E(T)$  isomorphically onto  $E(S)$ ;  $\psi$  is called a *covering homomorphism*. We show that any bountiful semigroup has an  $E$ -unitary cover on which  $\sigma \cap \mu = \iota$ . The latter property holds for all orthodox semigroups by [13, Lemma 2.2], and so our result is simply a generalisation of the existence of  $E$ -unitary orthodox covers for orthodox semigroups due independently to McAlister [13], Szendrei [22] and Takizawa [23]. As mentioned in the introduction, Simmons has established the existence of  $E$ -unitary covers for bountiful semigroups, but the covers he constructs are always infinite.

**Theorem 3.4.** *Let  $S$  be a bountiful semigroup. Then  $S$  has an  $E$ -unitary cover  $T$  such that  $\sigma \cap \mu = \iota$  on  $T$ , and  $T$  can be taken to be finite if  $S$  is finite.*

Our approach is inspired by that for orthodox semigroups in [13]. We start with the construction of  $T$ , and then in a series of lemmas show that  $T$  has the desired properties.

Let  $S$  be a bountiful semigroup. Then  $S/\delta$  is ample, and so by Theorem 3.2, it has a proper cover  $V$  with covering homomorphism  $\alpha$  say. Let

$$T = \{(s\mu, v) \in S/\mu \times V : v\alpha = s\delta\}.$$

It is easy to see that  $T$  is a subsemigroup of the direct product  $S/\mu \times V$ . We show, in a sequence of lemmas, that  $T$  is a cover of the required type.

**Lemma 3.5.** *The congruence  $\delta$  is idempotent pure.*

*Proof.* If  $s \in S$  is such that  $s\delta$  is idempotent, then since  $\delta$  is good, we have  $s\delta = h\delta$  for some  $h \in E(S)$  by Lemma 2.3. From the definition of  $\delta$ , we have  $s = ihj$  where  $i, j$  are idempotents. As  $E(S)$  is a subsemigroup, we see that  $s$  is idempotent.  $\square$

**Lemma 3.6.** *The idempotents of  $T$  form a subsemigroup.*

*Proof.* If  $e \in E(S)$  and  $v \in E(V)$  with  $v\alpha = e\delta$ , then clearly,  $(e\mu, v) \in E(T)$ .

On the other hand, if  $(s\mu, v) \in T$  is idempotent, then  $(s\mu)^2 = s\mu$  and  $v^2 = v$ . From the latter we have  $(s\delta)^2 = s\delta$  so that by Lemma 3.5,  $s$  is idempotent. It follows that

$$E(T) = \{(e\mu, v) \in S/\mu \times E(V) : e \in E(S) \text{ and } v\alpha = e\delta\}.$$

Since  $E(S)$  and  $E(V)$  are subsemigroups of  $S$  and  $V$  respectively, it follows that  $E(T)$  is a subsemigroup of  $T$ .  $\square$

**Lemma 3.7.** *Let  $(s\mu, u) \in T$ . If  $e \in E(S) \cap R_s^*$ , then  $(e\mu, u^+) \in T$ .*

*Proof.* Since  $(s\mu, u) \in T$ , we have  $s\delta = u\alpha$ . Now  $\alpha$  preserves  $\mathcal{R}^*$ , and so  $u^+\alpha\mathcal{R}^*u\alpha$ , that is,  $u^+\alpha\mathcal{R}^*s\delta$ . As  $\delta$  is also good and  $S/\delta$  is ample, this gives  $u^+\alpha = e\delta$  and the lemma follows.  $\square$

Next, we describe  $\mathcal{R}^*$  in  $T$ .

**Lemma 3.8.** *Let  $(A, u), (B, v) \in T$ . Then  $(A, u)\mathcal{R}^*(B, v)$  if and only if  $A\mathcal{R}^*B$  in  $S/\mu$  and  $u\mathcal{R}^*v$  in  $V$ .*

*Proof.* Let  $(A, u), (B, v) \in T$ , and suppose that  $A\mathcal{R}^*B$  and  $u\mathcal{R}^*v$ . If  $(X, x), (Y, y) \in T$  and  $(X, x)(A, u) = (Y, y)(A, u)$ , then  $XA = YA$  and  $xu = yu$  so that  $XB = YB$  and  $xv = yv$ , that is,  $(X, x)(B, v) = (Y, y)(B, v)$ . Similarly,  $(X, x)(A, u) = (A, u)$  implies  $(X, x)(B, v) = (B, v)$ . Together with the opposite implications, this gives  $(A, u)\mathcal{R}^*(B, v)$ .

Conversely, suppose that  $(A, u)\mathcal{R}^*(B, v)$  and let  $A = r\mu, B = s\mu$ . Now let  $e, f, u^+, v^+$  be idempotents in the  $\mathcal{R}^*$ -classes of  $r, s, u$  and  $v$  respectively. Put  $E = e\mu$  and  $F = f\mu$ . By Lemma 3.7,  $(E, u^+), (F, v^+) \in T$ , and, since  $\mu$  is good,  $A\mathcal{R}^*E$  and  $B\mathcal{R}^*F$ . Hence, by the first part,  $(A, u)\mathcal{R}^*(E, u^+)$  and  $(B, v)\mathcal{R}^*(F, v^+)$  so that  $(E, u^+)\mathcal{R}^*(F, v^+)$ . But these elements are idempotents, and so

$$(E, u^+)(F, v^+) = (F, v^+) \text{ and } (F, v^+)(E, u^+) = (E, u^+).$$

Comparing coordinates gives  $E\mathcal{R}F$  whence  $A\mathcal{R}^*B$ , and, since  $V$  is ample,  $u^+ = v^+$  so that  $u\mathcal{R}^*v$ .  $\square$

Notice that the second part of the proof shows that every element of  $T$  is  $\mathcal{R}^*$ -related to an idempotent. Similarly, each element is  $\mathcal{L}^*$ -related to an idempotent so that we have the following corollary.

**Corollary 3.9.**  *$T$  is abundant.*

To show that  $T$  is bountiful, we use the following characterisation of idempotent connected abundant semigroups [1, Lemma 2.3].

**Lemma 3.10.** *Let  $S$  be an abundant semigroup. Then  $S$  is IC if and only if it satisfies the following two conditions for all  $a \in S$ :*

- (1) *for some  $h \in E(S) \cap R_a^*$  and all  $f \leq h$ , there exists  $b \in S$  such that  $fa = ab$ ;*
- (2) *for some  $k \in E(S) \cap L_a^*$  and all  $e \leq k$ , there exists  $c \in S$  such that  $ae = ca$ .*

We note that the elements  $b$  and  $c$  can be taken to be idempotent, for if  $S$  is IC and  $B = \langle E(S) \rangle$ , then there is a connecting isomorphism  $\beta: \langle E(hBh) \rangle \rightarrow \langle E(kBk) \rangle$  and  $b$  and  $c$  may be chosen to be  $f\beta$  and  $e\beta^{-1}$  respectively. Furthermore, the discussion in [1] shows that the word ‘‘some’’ in conditions (1) and (2) of the lemma may be replaced by the word ‘‘all’’. The proof of the following lemma uses both versions.

**Lemma 3.11.**  *$T$  is bountiful.*

*Proof.* By Corollary 3.9 and Lemma 3.6,  $T$  is abundant and its idempotents form a sub-semigroup. All that remains is to show that  $T$  is IC.

Let  $(s\mu, v) \in T$  so that  $s\delta = v\alpha$ , and let  $h \in E(S)$  be  $\mathcal{R}^*$ -related to  $s$ . By Lemma 3.7,  $(h\mu, v^+) \in T$ , and by Lemma 3.8,  $(h\mu, v^+)\mathcal{R}^*(s\mu, v)$ . We show that condition (1) of Lemma 3.10 holds using  $(h\mu, v^+)$  as the idempotent in the  $\mathcal{R}^*$ -class of  $(s\mu, v)$ . First, we remark that since  $(h\mu, v^+) \in T$ , we have

$$h\delta = v^+\alpha. \tag{1}$$

Let  $(e\mu, f) \in E(T)$  where  $e \in E(S)$ ,  $f \in E(V)$  and  $f\alpha = e\delta$ . If  $(e\mu, f) \leq (h\mu, v^+)$ , then clearly  $e\mu \leq h\mu$  and  $f \leq v^+$ . Hence  $e\mu = (eh)\mu = (he)\mu$ , and so, using (1),

$$(eh)\delta = (e\delta)(h\delta) = (f\alpha)(v^+\alpha) = (fv^+)\alpha = f\alpha.$$

Similarly,  $(he)\delta = f\alpha$  so that  $e\delta = (eh)\delta = (he)\delta$ . As  $\mu \cap \delta = \iota$ , we obtain  $e = eh = he$ , that is,  $e \leq h$ .

Now  $S$  is bountiful, so by Lemma 3.10 and the remarks following it, there is an idempotent  $k \in S$  with  $es = sk$ .

Since  $V$  is a proper cover of  $S/\delta$ , there is an idempotent  $u$  in  $V$  with  $u\alpha = k\delta$  so that  $(k\mu, u) \in E(T)$ .

Now

$$(fv)\alpha = (f\alpha)(v\alpha) = (e\delta)(s\delta) = (es)\delta = (sk)\delta = (s\delta)(k\delta) = (v\alpha)(u\alpha) = (vu)\alpha.$$

As  $\alpha$  preserves  $*$  and is one-one on  $E(V)$ , we get  $(fv)^* = (vu)^* = v^*u$ . Hence, since  $V$  is ample,

$$vu = vv^*u = v(fv)^* = fv.$$

Now we have  $(k\mu, u) \in T$  and

$$(e\mu, f)(s\mu, v) = ((es)\mu, fv) = ((sk)\mu, vu) = (s\mu, v)(k\mu, u),$$

and so condition (1) of Lemma 3.10 holds for  $T$ . Similarly, condition (2) also holds, and so by the lemma,  $T$  is IC, and hence bountiful.  $\square$

**Lemma 3.12.**  *$T$  is  $E$ -unitary.*

*Proof.* Let  $(s\mu, v) \in T$  so that  $s\delta = v\alpha$ , and let  $(e\mu, f) \in E(T)$  where  $e \in E(S)$  and  $e\delta = f\alpha$ . Suppose that  $(s\mu, v)(e\mu, f) \in E(T)$ . Then  $vf \in E(V)$  and so  $v \in E(V)$  since  $V$  is proper and hence  $E$ -unitary.

Also  $s\delta = v\alpha$  is an idempotent of  $S/\delta$  and so, by Lemma 3.5,  $s$  is idempotent. Thus  $(s\mu, v) \in E(T)$  and it follows that  $T$  is  $E$ -unitary.  $\square$

We will find the following lemma useful; it is essentially part of Proposition 2.1 of [2].

**Lemma 3.13.** *Let  $a, b$  be elements of an abundant semigroup  $S$ . Then the following are equivalent:*

- (1)  $a\mu_s b$ ,
- (2)  $ae\mathcal{R}^*be$  and  $ea\mathcal{L}^*eb$  for all  $e \in E(S)$ .

**Lemma 3.14.** *If  $(r\mu, u), (s\mu, v) \in T$  and  $((r\mu, u), (s\mu, v)) \in \sigma \cap \mu$ , then  $(r\mu, u) = (s\mu, v)$ .*

*Proof.* Let  $e \in E(S)$  and  $f$  be any idempotent in  $V$  such that  $(e\mu, f) \in E(T)$ . Then, by Lemma 3.13 we have

$$(r\mu, u)(e\mu, f)\mathcal{R}^*(s\mu, v)(e\mu, f) \text{ and } (e\mu, f)(r\mu, u)\mathcal{L}^*(e\mu, f)(s\mu, v).$$

By Lemma 3.8 and its dual,  $(r\mu)(e\mu)\mathcal{R}^*(s\mu)(e\mu)$  and  $(e\mu)(r\mu)\mathcal{L}^*(e\mu)(s\mu)$ . It follows from Lemma 2.3 that every idempotent of  $S/\mu$  is of the form  $e\mu$  for an idempotent  $e$  of  $S$ , and so, by Lemma 3.13 again, we have  $(r\mu, s\mu) \in \mu_{S/\mu}$ . Hence by Theorem 2.5,  $\mu_{S/\mu} = \iota$  so that  $r\mu = s\mu$ .

Now  $\mu \subseteq \mathcal{R}^*$ , so that  $(r\mu, u)\mathcal{R}^*(s\mu, v)$ . By Lemma 3.8, we have  $u\mathcal{R}^*v$ . As  $(r\mu, u)\sigma(s\mu, v)$ , we have  $(e\mu, f)(r\mu, u) = (s\mu, v)(h\mu, k)$  for some idempotents  $(e\mu, f)$  and  $(h\mu, k)$  of  $T$ , so that  $fu = vk$ , and, since  $V$  is ample, it follows that  $u\sigma v$ . Now  $(u, v) \in \mathcal{R}^* \cap \sigma$  and  $V$  is proper, so  $u = v$ . Hence  $(r\mu, u) = (s\mu, v)$ .  $\square$



The proof of Theorem 3.4 is completed by the next lemma.

**Lemma 3.15.** *The mapping  $\theta : T \rightarrow S$  given by  $(s\mu, v)\theta = s$  is a well defined good homomorphism onto  $S$  which maps  $E(T)$  isomorphically onto  $E(S)$ .*

*Proof.* First, we note that if  $(r\mu, u) = (s\mu, v)$ , then  $r\delta = u\alpha = v\alpha = s\delta$  so that  $(r, s) \in \mu \cap \delta$ , and so, by the remarks preceding Proposition 3.3,  $r = s$ . Thus  $\theta$  is well defined. It is clear that  $\theta$  is a homomorphism. If  $s \in S$ , then, since  $\alpha$  is surjective, there is an element  $v \in V$  such that  $v\alpha = s\delta$ , so that  $(s\mu, v) \in T$  and  $\theta$  is surjective; this also shows that  $\theta$  maps  $E(T)$  onto  $E(S)$ .

To see that  $\theta$  is one-one on  $E(T)$ , let  $(e\mu, v), (f\mu, w) \in E(T)$  and suppose that  $(e\mu, v)\theta = (f\mu, w)\theta$ . Then  $e = f$  so that certainly  $e\mu = f\mu$ . Also,  $v, w \in E(V)$  and  $v\alpha = e\delta = f\delta = w\alpha$  so that  $v = w$  since  $\alpha$  is idempotent separating.

Finally, to see that  $\theta$  is good, suppose that  $(r\mu, u), (s\mu, v) \in T$  with  $(r\mu, u)\mathcal{R}^*(s\mu, v)$ . By Lemma 3.8,  $r\mu\mathcal{R}^*s\mu$  and  $u\mathcal{R}^*v$ . From the latter, we get  $r\delta\mathcal{R}^*s\delta$  since  $r\delta = u\alpha$ ,  $s\delta = v\alpha$  and  $\alpha$  is good. Let  $e, f \in E(S)$  be such that  $e\mathcal{R}^*r$  and  $f\mathcal{R}^*s$ . Then  $e\mu\mathcal{R}^*f\mu$  and  $e\delta\mathcal{R}^*f\delta$ . It follows from this that  $(ef, f), (fe, e) \in \mu \cap \delta$ . Hence  $ef = f$  and  $fe = e$  so that  $e\mathcal{R}^*f$  and thus  $r\mathcal{R}^*s$  as desired.

Similarly,  $\theta$  preserves  $\mathcal{L}^*$  and so  $\theta$  is good.  $\square$

Having proved Theorem 3.4, it is now easy to prove the following result which completes the proof of Theorem 1.

**Proposition 3.16.** *If  $S$  is a finite bountiful semigroup with  $\mathcal{H} \subseteq \mu$ , then  $S \in \mathbf{A} \vee \mathbf{G}$ .*

*Proof.* Let  $T$  be a finite  $E$ -unitary cover of  $S$  with  $\mu \cap \sigma = \iota$  and covering map  $\theta : T \rightarrow S$ . Now  $\mu \cap \sigma = \iota$ , and so  $T$  can be embedded (as a subdirect product) in  $T/\mu \times T/\sigma$ . Since  $T/\sigma$  is cancellative and finite, we have  $T/\sigma \in \mathbf{G}$ .

Now  $\mathcal{H} \subseteq \mu$  on  $S$ , so  $S/\mu$  has only trivial subgroups, and hence  $S/\mu \in \mathbf{A}$ .

We claim that  $T/\mu \cong S/\mu$ . To see this, it is enough to show that for  $a, b \in T$  we have  $a\mu_T b$  if and only if  $(a\theta)\mu_S(b\theta)$ . Using Lemma 3.13 and the fact that  $\theta$  is good, it is easy to see that  $a\mu_T b$  implies  $(a\theta)\mu_S(b\theta)$ .

Conversely, if  $(a\theta)\mu_S(b\theta)$ , then, again by Lemma 3.13, we have that  $(ae)\theta\mathcal{R}^*(be)\theta$  for all  $e \in E(T)$ . Let  $f, h \in E(T)$  be in the  $\mathcal{R}^*$ -classes of  $ae$  and  $be$  respectively. Since  $\theta$  is good, we obtain  $f\theta\mathcal{R}^*h\theta$ . But these elements are idempotent, so  $(fh)\theta = (f\theta)(h\theta) = h\theta$  and  $(hf)\theta = f\theta$ . Now  $T$  is bountiful, so  $fh$  and  $hf$  are idempotents, and so  $fh = h$  and  $hf = f$  since  $\theta$  is idempotent separating. Hence  $f\mathcal{R}h$  and so  $ae\mathcal{R}^*be$ . Similarly,  $ea\mathcal{L}^*eb$  for all  $e \in E(T)$  so that  $a\mu_T b$ .

Hence the claim is proved, so that  $T/\mu \in \mathbf{A}$ , and thus  $S \in \mathbf{A} \vee \mathbf{G}$ , since  $S$  divides  $T/\mu \times T/\sigma$ .  $\square$

## REFERENCES

- [1] S. Armstrong, Structure of concordant semigroups, *J. Algebra* **118** (1988), 205–260.
- [2] A. El Qallali and J. B. Fountain, Idempotent-connected abundant semigroups, *Proc. Roy. Soc. Edinburgh* **91A** (1981), 79–90.

- [3] A. El Qallali and J. B. Fountain, Quasi-adequate semigroups, *Proc. Royal Soc. Edinburgh* **91A** (1981), 91–99.
- [4] J. Fountain, A class of right PP monoids, *Quart. J. Math. Oxford* **28** (1977), 285–300.
- [5] J. Fountain, Abundant semigroups, *Proc. London Math. Soc.*, **44** (1982), 103–129.
- [6] J. Fountain and G. M. S. Gomes, Proper covers of ample semigroups, *Proc. Edinb. Math. Soc.*, to appear.
- [7] Xiaojiang Guo, F-abundant semigroups, *Glasgow Math. J.* **43** (2001), 153–163.
- [8] T. E. Hall, On regular semigroups, *J. Algebra* **24** (1973), 1–24.
- [9] J. M. Howie, *Fundamentals of semigroup theory*, Oxford University Press, 1995.
- [10] M.V. Lawson, *The Structure Theory of Abundant Semigroups*, D.Phil. thesis, York, 1985.
- [11] M. V. Lawson, The structure of type A semigroups, *Quart. J. Math. Oxford* **37** (1986), 279–298.
- [12] D. B. McAlister, One-to-one partial right right translations of a right cancellative semigroup. *J. Algebra* **43** (1976), 231–251.
- [13] D. B. McAlister, On a problem of M. P. Schützenberger, *Proc. Edinb. Math. Soc.* **23** (1980), 243–247.
- [14] D. B. McAlister, Regular semigroups, fundamental semigroups and groups, *J. Austral. Math. Soc. Ser. A* **29** (1980), 475–503.
- [15] K. S. S. Nambooripad, *Structure of regular semigroups I*, Mem. Amer. Math. Soc. **22** (1979), no. 224.
- [16] F. Pastijn, A representation of a semigroup by a semigroup of matrices over a group with zero, *Semigroup Forum* **10** (1975), 238–239.
- [17] J.-E. Pin, *Varieties of Formal Languages*, North Oxford Academic, 1986.
- [18] J. Rhodes and B. Tilson, Improved lower bounds for the complexity of finite semigroups, *J. Pure Appl. Algebra* **2** (1972), 13–71.
- [19] Zhang Ronghua, private communication (1996).
- [20] C. P. Simmons, On abundant  $E$ -monoids, preprint 2002.
- [21] B. Steinberg, private communication (2002).
- [22] M. B. Szendrei, On a pull-back diagram for orthodox semigroups, *Semigroup Forum* **20** (1980), 1–10.
- [23] K. Takizawa, Orthodox semigroups and  $E$ -unitary regular semigroups, *Bull. Tokyo Gakugei Univ., Series IV* **31** (1979), 41–43.