



Full site index  
Go

Full search  
Search

Advertiser

**Latest News**

- Florida justices ponder recount
- Latest controversy takes familiar path-to court
- Officials urged to give military vote 2nd look
- Freed man awaits next step
- Israel attacks Gaza following school bus attack
- President's resignation creates turmoil in Peru
- Japanese Parliament fails to oust prime minister

[More Latest News](#)

**Markets**

*DOW:*  
-167.27 10462.6  
*NASDAQ:*  
-151.55 2875.64  
*S&P 500:*  
-25.1 1342.62

**Find a quote:**

**Search**

[Full market report](#)

**Web specials**



Special report:  
Gateway to Gridlock

## CHIPS MAY TAKE QUANTUM LEAP SCIENTISTS EXPLORE MANUFACTURING EVEN THINNER TRANSISTORS

**By Ian Austen**  
New York Times News Service  
*November 13, 2000*

Microchips are becoming more sophisticated and faster largely because their designers keep coming up with ways to make the channels that are etched to create transistors increasingly tiny. But a day of reckoning may arrive. If transistors reach a size of 25 nanometers (roughly the width of 250 hydrogen atoms set side by side) the electrons flowing through them would probably no longer play by the rules of classical physics.

Instead, the sometimes bizarre effects described by quantum theory would take over.

The good news for Intel and other chipmakers is that they still have a long way to go before their chips start misbehaving. The most advanced chips widely available at the moment have transistors that are no thinner than 180 to 220 nanometers, roughly 2,000 atoms wide. There is concern, however, that the limitations of chipmaking technology may prevent the industry from creating circuitry thinner than that.

Now, in something of a reversal, a paper in the journal *Physical Review Letters* suggests that quantum theory may be a chip designer's friend, at least when it comes to manufacturing. In it, four quantum physics researchers from the Jet Propulsion Laboratory, managed for NASA by the California Institute of Technology, and two scientists from the University of Wales suggest that by embracing quantum theory, chipmakers may find a way to shrink transistors well beyond current limits while avoiding the cost of building entirely new factories to do it.

"It's a whole new way of looking at chipmaking," said Daniel Abrams, a research scientist at the Jet Propulsion Laboratory and a co-author of the research report.

**Well Read.**  
**Books**  
chicagotribune.com

*Advertising inform*

Gateway to Gridlock

e-tailing

Illinois school report card

Investigative report:  
Forbidden friendships  
between cops and  
criminals

Dangerous care: Nurses'  
hidden role in medical  
error

1999 Illinois State Police  
crime report

Jelly Roll blues

---

*More specials*

At its most basic level, chip manufacturing today is similar to what goes on when you take your film to a camera shop for processing and printing. Light is beamed onto a photosensitive surface through a mask, a chip-design template that acts much like a negative. The resulting pattern is then etched in place to form the transistors in a process called optical lithography.

As transistors get smaller, chipmakers need to use light of increasingly smaller wavelengths to make their exposures.

The most common approach uses optics to create an effect called diffraction. Using diffraction, waves of deep ultraviolet light are generated by lasers and then passed through optical systems to decrease their wavelength even further. But the current 180- to 220-nanometer light waves used today will not be able to make transistors smaller than 124 nanometers.

A number of proposals for generating smaller beams are in various stages of development. One uses ultraviolet light with short wavelengths, another X-rays and a third beams of electrons. None of the techniques, however, will work with current chip lithography systems, and most of them may prove too slow and costly for large-scale manufacturing. The recent quantum physics paper may give chipmakers a way to adapt their current factories to generate tiny wavelengths of light that can generate very tiny circuitry.

At the heart of the technique is what Abrams calls "one of the weird quantum effects." Photons--particles of light--usually do not interact with one another. But in some cases two or more photons can become, in quantum theory terms, entangled, linking their fates and properties together in a strange way.

"They're correlated in a funny way so that they affect each other even at a distance," Abrams said. "Einstein called it 'spooky action at a distance.'"

The process suggested in the paper begins by zapping crystals made of either potassium diphosphate and potassium triphosphate with small-wavelength laser light. The crystals, in turn, spew out a stream of entangled photons.

The entangled photons are then aimed at two slits. When light of various types is aimed at a diffraction grating, the light passes through the slits and forms characteristic patterns on the other side. When an entangled pair of photons is squeezed through the slits, Abrams said, they appear to have twice the energy of ordinary photons making the same trip. (And, in another oddity of quantum theory, the entangled pairs pass through both slits simultaneously.)

When the entangled pairs recombine on the far side of the slit, they form a wavelength that is half the size of what can be generated by using unentangled photons. With the technology, light from a conventional 248-nanometer laser could be reduced to a wavelength of 62 nanometers to make transistors of that size, about a third of what can be achieved with current methods.

The potential for miniaturization does not stop there.

Entangling three or even more photons and then passing them through the slit should, in theory, result in even shorter wavelengths of light, Abrams said, and that should allow chipmakers to create even smaller circuitry "while still using a regular laser and optics."

But that "cute trick" may take years to make its way into chip plants, Abrams cautioned. "To go from a theoretical idea to production is a large step," he said.

There are some doubters. Paul Kwiat, a physicist at the Los Alamos National Laboratory, said that he did not dispute any of the science in the recent paper. But he warned that the proposed technique might encounter several insurmountable problems on the factory floor. "We don't have really good sources for bright, entangled photons."

The crystal method generated about 1 million entangled photons per second, Kwiat said. "That may sound impressive, but even a simple laser pointer creates 1 billion million."

So, he said, it may be hard to produce enough entangled photons for practical use. He also doubts that any practical method for actually aiming the entangled photons can be found, and that would hinder the adoption of the new strategy.

"I'm a bit pessimistic for its widespread application," Kwiat said.

[E-mail this story to a friend](#)

[More articles on the Business News home page](#)

[News](#) | [Sports](#) | [Business](#) | [Tech](#) | [Leisure](#) | [Travel](#) | [Classified: Jobs · Cars · Homes](#)

[Top of page](#) | [Home](#) | [Copyright and terms of service](#) | [Privacy policy](#) | [Subscribe](#) | [Customer service](#) | [Advertise](#)

**YOUR SCHOOL**

**ChicagoSports.com**  
TRY TO KEEP UP.