

Weakly U -abundant semigroups

Yanhui Wang


The University of York, UK

9th Nov. 2011

NBSAN

Outline: U a biordered set


Weakly U -abundant Semigps With (C)



Groupoids

Definition A *groupoid* is a category G in which for every $p \in \text{Mor}G$ we have $p^{-1} \in \text{Mor}G$ with

$$pp^{-1} = l_{d(p)} \text{ and } p^{-1}p = l_{r(p)}.$$



Inverse semigroups \rightarrow Groupoids

Let S be an inverse semigroup with semilattice of idempotents E . We construct a groupoid $C(S)$ as follows:

$$\text{Ob } C(S) = E, \text{ Mor } C(S) = S, \mathbf{d}(x) = xx', \mathbf{r}(x) = x'x$$

and a partial binary operation \cdot is defined by the rule that for any $x, y \in S$,

$$x \cdot y = \begin{cases} xy & \text{if } \mathbf{r}(x) = \mathbf{d}(y), \\ \text{undefined,} & \text{otherwise.} \end{cases}$$

where xy is the product of x and y in S .

Let $G = (G, \cdot)$ be a small groupoid. Let S be the semigroup obtained from G by declaring all undefined products to be 0.

Fact 1. $S = G \cup \{0\}$ is an inverse semigroup with 0 with $E(S) = E(G) \cup \{0\}$.

Let $G = (G, \cdot)$ be a small groupoid. Let S be the semigroup obtained from G by declaring all undefined products to be 0.

Fact 1. $S = G \cup \{0\}$ is an inverse semigroup with 0 with $E(S) = E(G) \cup \{0\}$.

Inductive groupoids

Let $G = (G, \cdot)$ be a small groupoid with $E(G) = E$. Suppose that \leq is a partial order on G . Suppose also

- 1 $x \leq y$ implies that $x^{-1} \leq y^{-1}$;
 - 2 $x \leq y, u \leq v, \exists x \cdot u, \exists y \cdot v$ implies that $x \cdot u \leq y \cdot v$;
 - 3 if $a \in G$ and $e \in E$ with $e \leq \mathbf{d}(a)$, then there exists a unique **restriction** $(e|a) \in G$ with $\mathbf{d}(e|a) = e$ and $(e|a) \leq a$;
 - 4 if $a \in G$ and $e \in E$ with $e \leq \mathbf{r}(a)$, then there exists a unique **co-restriction** $(a|e) \in G$ with $\mathbf{r}(a|e) = e$ and $(a|e) \leq a$;
- then $G = (G, \cdot, \leq)$ is called an **ordered groupoid**. If in addition
- 5 E is a semilattice.

Then $G = (G, \cdot, \leq)$ is called an **inductive groupoid**.

Inductive groupoids

Let $G = (G, \cdot)$ be a small groupoid with $E(G) = E$. Suppose that \leq is a partial order on G . Suppose also

- 1 $x \leq y$ implies that $x^{-1} \leq y^{-1}$;
 - 2 $x \leq y, u \leq v, \exists x \cdot u, \exists y \cdot v$ implies that $x \cdot u \leq y \cdot v$;
 - 3 if $a \in G$ and $e \in E$ with $e \leq \mathbf{d}(a)$, then there exists a unique **restriction** $(e|a) \in G$ with $\mathbf{d}(e|a) = e$ and $(e|a) \leq a$;
 - 4 if $a \in G$ and $e \in E$ with $e \leq \mathbf{r}(a)$, then there exists a unique **co-restriction** $(a|e) \in G$ with $\mathbf{r}(a|e) = e$ and $(a|e) \leq a$;
- then $G = (G, \cdot, \leq)$ is called an **ordered groupoid**. If in addition
- 5 E is a semilattice.

Then $G = (G, \cdot, \leq)$ is called an **inductive groupoid**.

Inductive groupoids

Let $G = (G, \cdot)$ be a small groupoid with $E(G) = E$. Suppose that \leq is a partial order on G . Suppose also

- 1 $x \leq y$ implies that $x^{-1} \leq y^{-1}$;
 - 2 $x \leq y, u \leq v, \exists x \cdot u, \exists y \cdot v$ implies that $x \cdot u \leq y \cdot v$;
 - 3 if $a \in G$ and $e \in E$ with $e \leq \mathbf{d}(a)$, then there exists a unique **restriction** $(e|a) \in G$ with $\mathbf{d}(e|a) = e$ and $(e|a) \leq a$;
 - 4 if $a \in G$ and $e \in E$ with $e \leq \mathbf{r}(a)$, then there exists a unique **co-restriction** $(a|e) \in G$ with $\mathbf{r}(a|e) = e$ and $(a|e) \leq a$;
- then $G = (G, \cdot, \leq)$ is called an **ordered groupoid**. If in addition
- 5 E is a semilattice.

Then $G = (G, \cdot, \leq)$ is called an **inductive groupoid**.

Inductive groupoids

Let $G = (G, \cdot)$ be a small groupoid with $E(G) = E$. Suppose that \leq is a partial order on G . Suppose also

- 1 $x \leq y$ implies that $x^{-1} \leq y^{-1}$;
- 2 $x \leq y, u \leq v, \exists x \cdot u, \exists y \cdot v$ implies that $x \cdot u \leq y \cdot v$;
- 3 if $a \in G$ and $e \in E$ with $e \leq \mathbf{d}(a)$, then there exists a unique **restriction** $(e|a) \in G$ with $\mathbf{d}(e|a) = e$ and $(e|a) \leq a$;
- 4 if $a \in G$ and $e \in E$ with $e \leq \mathbf{r}(a)$, then there exists a unique **co-restriction** $(a|e) \in G$ with $\mathbf{r}(a|e) = e$ and $(a|e) \leq a$;
then $G = (G, \cdot, \leq)$ is called an **ordered groupoid**. If in addition
- 5 E is a semilattice.

Then $G = (G, \cdot, \leq)$ is called an **inductive groupoid**.

Inverse semigroups \rightarrow Inductive groupoids

Let S be an inverse semigroup with semilattice of idempotents E . Let $C(S) = (S, \cdot)$ be defined as above.

Partial Order: For any $a, b \in S$, $a \leq b \Leftrightarrow a = eb$ for some $e \in E$

Fact 2. $C(S) = (S, \cdot, \leq)$ is an inductive groupoid with

$$(e|a) = ea, (a|e) = ae.$$

Inverse semigroups \rightarrow Inductive groupoids

Let S be an inverse semigroup with semilattice of idempotents E . Let $C(S) = (S, \cdot)$ be defined as above.

Partial Order: For any $a, b \in S$, $a \leq b \Leftrightarrow a = eb$ for some $e \in E$

Fact 2. $C(S) = (S, \cdot, \leq)$ is an inductive groupoid with

$$(e|a) = ea, (a|e) = ae.$$

Inverse semigroups \rightarrow Inductive groupoids

Let S be an inverse semigroup with semilattice of idempotents E . Let $C(S) = (S, \cdot)$ be defined as above.

Partial Order: For any $a, b \in S$, $a \leq b \Leftrightarrow a = eb$ for some $e \in E$

Fact 2. $C(S) = (S, \cdot, \leq)$ is an inductive groupoid with

$$(e|a) = ea, (a|e) = ae.$$

Inductive groupoids \rightarrow Inverse semigroups

Let $G = (G, \cdot, \leq)$ be an inductive groupoid.

We define a *pseudo-product* \otimes on G by the rule that

$$a \otimes b = (a|\mathbf{r}(a) \wedge \mathbf{d}(b)) \cdot (\mathbf{r}(a) \wedge \mathbf{d}(b)|b).$$

Fact 3. $\mathcal{S}(G) = (G, \otimes)$ is an inverse semigroup (having the same partial order as G).

Ehresmann-Schein-Nambooripad

The category \mathbf{I} of inverse semigroups and morphisms is isomorphic to the category \mathbf{G} of **inductive groupoids** and inductive functors.



Inductive groupoids \rightarrow Inverse semigroups

Let $G = (G, \cdot, \leq)$ be an inductive groupoid.

We define a *pseudo-product* \otimes on G by the rule that

$$a \otimes b = (a|r(a) \wedge \mathbf{d}(b)) \cdot (r(a) \wedge \mathbf{d}(b)|b).$$

Fact 3. $\mathcal{S}(G) = (G, \otimes)$ is an inverse semigroup (having the same partial order as G).

Ehresmann-Schein-Nambooripad

The category \mathbf{I} of inverse semigroups and morphisms is isomorphic to the category \mathbf{G} of *inductive groupoids* and inductive functors.



Inductive groupoids \rightarrow Inverse semigroups

Let $G = (G, \cdot, \leq)$ be an inductive groupoid.

We define a *pseudo-product* \otimes on G by the rule that

$$a \otimes b = (a|r(a) \wedge \mathbf{d}(b)) \cdot (r(a) \wedge \mathbf{d}(b)|b).$$

Fact 3. $\mathcal{S}(G) = (G, \otimes)$ is an inverse semigroup (having the same partial order as G).

Ehresmann-Schein-Nambooripad

The category \mathbf{I} of inverse semigroups and morphisms is isomorphic to the category \mathbf{G} of [inductive groupoids](#) and inductive functors.



Green's star equivalences


Definition

- Given a semigroup S , for any $a, b \in S$,
 $a \mathcal{L}^* b$ if $a \mathcal{L} b$ in a semigroup T such that $S \subseteq T$.

Dually, The relation \mathcal{R}^* on S is defined.

$$\mathcal{D}^* = \mathcal{L}^* \vee \mathcal{R}^* .$$

$$\mathcal{H}^* = \mathcal{L}^* \wedge \mathcal{R}^* .$$



$$\mathcal{L} \subseteq \mathcal{L}^*$$

$$\mathcal{R} \subseteq \mathcal{R}^*$$

- A semigroup is called U -abundant if each \mathcal{L}^* -class and each \mathcal{R}^* -class contains an idempotent.

Green's star equivalences


Definition

- Given a semigroup S , for any $a, b \in S$,
 $a \mathcal{L}^* b$ if $a \mathcal{L} b$ in a semigroup T such that $S \subseteq T$.

Dually, The relation \mathcal{R}^* on S is defined.

$$\mathcal{D}^* = \mathcal{L}^* \vee \mathcal{R}^* .$$

$$\mathcal{H}^* = \mathcal{L}^* \wedge \mathcal{R}^* .$$



$$\mathcal{L} \subseteq \mathcal{L}^*$$

$$\mathcal{R} \subseteq \mathcal{R}^*$$

- A semigroup is called **U -abundant** if each \mathcal{L}^* -class and each \mathcal{R}^* -class contains an idempotent.

(IC) and concordant semigroups

Let S be an abundant semigroup with $E(S) = U$. If $a \in S$, then a^* and a^\dagger denote typical idempotents in L_a^* and R_a^* , respectively.

Then S is idempotent-connected (IC) if for any $a \in S$ and for some a^\dagger , a^* , there exists a bijection $\alpha : \langle a^\dagger \rangle \rightarrow \langle a^* \rangle$ satisfying $xa = a(x\alpha)$ for all $x \in \langle a^\dagger \rangle$, where for any $e \in U$, $\langle e \rangle$ is the subsemigroup generated by idempotents of eUe .

An abundant semigroup S is called a **concordant semigroup** if S satisfies Condition (IC) and its set of idempotents generates a regular subsemigroup.



(IC) and concordant semigroups

Let S be an abundant semigroup with $E(S) = U$. If $a \in S$, then a^* and a^\dagger denote typical idempotents in L_a^* and R_a^* , respectively.

Then S is idempotent-connected (IC) if for any $a \in S$ and for some a^\dagger , a^* , there exists a bijection $\alpha : \langle a^\dagger \rangle \rightarrow \langle a^* \rangle$ satisfying $xa = a(x\alpha)$ for all $x \in \langle a^\dagger \rangle$, where for any $e \in U$, $\langle e \rangle$ is the subsemigroup generated by idempotents of eUe .

An abundant semigroup S is called a **concordant semigroup** if S satisfies Condition (IC) and its set of idempotents generates a regular subsemigroup.



(IC) and concordant semigroups

Let S be an abundant semigroup with $E(S) = U$. If $a \in S$, then a^* and a^\dagger denote typical idempotents in L_a^* and R_a^* , respectively.

Then S is idempotent-connected (IC) if for any $a \in S$ and for some a^\dagger , a^* , there exists a bijection $\alpha : \langle a^\dagger \rangle \rightarrow \langle a^* \rangle$ satisfying $xa = a(x\alpha)$ for all $x \in \langle a^\dagger \rangle$, where for any $e \in U$, $\langle e \rangle$ is the subsemigroup generated by idempotents of eUe .

An abundant semigroup S is called a **concordant semigroup** if S satisfies Condition (IC) and its set of idempotents generates a regular subsemigroup.



(IC) and concordant semigroups

Let S be an abundant semigroup with $E(S) = U$. If $a \in S$, then a^* and a^\dagger denote typical idempotents in L_a^* and R_a^* , respectively.

Then S is idempotent-connected (IC) if for any $a \in S$ and for some a^\dagger , a^* , there exists a bijection $\alpha : \langle a^\dagger \rangle \rightarrow \langle a^* \rangle$ satisfying $xa = a(x\alpha)$ for all $x \in \langle a^\dagger \rangle$, where for any $e \in U$, $\langle e \rangle$ is the subsemigroup generated by idempotents of eUe .

An abundant semigroup S is called a **concordant semigroup** if S satisfies Condition (IC) and its set of idempotents generates a regular subsemigroup.



Concordant semigroups \rightarrow Categories

Let S be a concordant semigroup with set of idempotents U .

Set

$$\mathbf{K}(S) = \{(e, a, f) : f \in L_a^* \cap U, e \in R_a^* \cap U\}.$$

We define a partial binary operation on $K(S)$ by

$$(e, a, f) \cdot (g, b, h) = \begin{cases} (e, ab, h) & \text{if } f = g \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Partial Order:

$(e, a, f) \leq (g, b, h) \Leftrightarrow e \omega g, a = eb$ and $f = e\beta$, where $\beta : \langle e \rangle \rightarrow \langle f \rangle$.

Fact 4: The set $(\mathbf{K}(S), \cdot, \leq)$ forms an inductive cancellative category.

Concordant semigroups \rightarrow Categories

Let S be a concordant semigroup with set of idempotents U .

Set

$$\mathbf{K}(S) = \{(e, a, f) : f \in L_a^* \cap U, e \in R_a^* \cap U\}.$$

We define a partial binary operation on $K(S)$ by

$$(e, a, f) \cdot (g, b, h) = \begin{cases} (e, ab, h) & \text{if } f = g \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Partial Order:

$(e, a, f) \leq (g, b, h) \Leftrightarrow e \omega g, a = eb$ and $f = e\beta$, where $\beta : \langle e \rangle \rightarrow \langle f \rangle$.

Fact 4: The set $(\mathbf{K}(S), \cdot, \leq)$ forms an inductive cancellative category.

Concordant semigroups \rightarrow Categories

Let S be a concordant semigroup with set of idempotents U .

Set

$$\mathbf{K}(S) = \{(e, a, f) : f \in L_a^* \cap U, e \in R_a^* \cap U\}.$$

We define a partial binary operation on $K(S)$ by

$$(e, a, f) \cdot (g, b, h) = \begin{cases} (e, ab, h) & \text{if } f = g \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Partial Order:

$(e, a, f) \leq (g, b, h) \Leftrightarrow e \omega g, a = eb$ and $f = e\beta$, where $\beta : \langle e \rangle \rightarrow \langle f \rangle$.

Fact 4: The set $(\mathbf{K}(S), \cdot, \leq)$ forms an inductive cancellative category.

Categories \rightarrow Concordant semigroups

Let (C, \cdot, \leq) be an inductive cancellative category with regular biordered set of objects E and ρ be a congruence on (C, \cdot, \leq) .

For any $x \in C$, \bar{x} denotes the ρ -class containing x . Define

$$\bar{x} \circ \bar{y} = \overline{(x \otimes y)_h},$$

where $h \in S(\mathbf{r}(x), \mathbf{d}(y))$.

Fact 5: The set $(C/\rho, \circ)$ forms a concordant semigroup.

Theorem: The category of inductive cancellative categories is equivalent to the category of concordant semigroups. (S. Armstrong, 1988)



Let (C, \cdot, \leq) be an inductive cancellative category with regular biordered set of objects E and ρ be a congruence on (C, \cdot, \leq) .

For any $x \in C$, \bar{x} denotes the ρ -class containing x . Define

$$\bar{x} \circ \bar{y} = \overline{(x \otimes y)_h},$$

where $h \in S(\mathbf{r}(x), \mathbf{d}(y))$.

Fact 5: The set $(C/\rho, \circ)$ forms a concordant semigroup.

Theorem: The category of inductive cancellative categories is equivalent to the category of concordant semigroups. (S. Armstrong, 1988)



Let (C, \cdot, \leq) be an inductive cancellative category with regular biordered set of objects E and ρ be a congruence on (C, \cdot, \leq) .

For any $x \in C$, \bar{x} denotes the ρ -class containing x . Define

$$\bar{x} \circ \bar{y} = \overline{(x \otimes y)_h},$$

where $h \in S(\mathbf{r}(x), \mathbf{d}(y))$.

Fact 5: The set $(C/\rho, \circ)$ forms a concordant semigroup.

Theorem: The category of inductive cancellative categories is equivalent to the category of concordant semigroups. (S. Armstrong, 1988)



$\tilde{\mathcal{L}}_U, \tilde{\mathcal{R}}_U$ equivalences

Definition

- Let $U \subseteq E(S)$. For any $a, b \in S$,
 $a \tilde{\mathcal{L}}_U b \Leftrightarrow (\forall e \in U)(ae = a \text{ if and only if } be = b)$,
 $a \tilde{\mathcal{R}}_U b \Leftrightarrow (\forall e \in U)(ea = a \text{ if and only if } eb = b)$,
 $\tilde{\mathcal{H}}_U = \tilde{\mathcal{L}}_U \wedge \tilde{\mathcal{R}}_U, \quad \tilde{\mathcal{D}}_U = \tilde{\mathcal{L}}_U \vee \tilde{\mathcal{R}}_U.$
- A semigroup S with $U \subseteq E(S)$ is said to be **weakly U -abundant** if each $\tilde{\mathcal{L}}_U$ -class and each $\tilde{\mathcal{R}}_U$ -class contains an idempotent in U .
- A weakly U -abundant semigroup S satisfies **Congruence Condition (C)** if $\tilde{\mathcal{L}}_U$ is a right congruence and $\tilde{\mathcal{R}}_U$ is a left congruence.
- A weakly E -abundant semigroup (S) is an **Ehresmann semigroup** if S satisfies (C) and E is a semilattice.
- A **weakly U -concordant semigroup** is a weakly U -abundant semigroup with (C) and $\langle U \rangle$ being a regular semigroup whose set of idempotents is U .



$\tilde{\mathcal{L}}_U, \tilde{\mathcal{R}}_U$ equivalences

Definition

- Let $U \subseteq E(S)$. For any $a, b \in S$,
 $a \tilde{\mathcal{L}}_U b \Leftrightarrow (\forall e \in U)(ae = a \text{ if and only if } be = b)$,
 $a \tilde{\mathcal{R}}_U b \Leftrightarrow (\forall e \in U)(ea = a \text{ if and only if } eb = b)$,
 $\tilde{\mathcal{H}}_U = \tilde{\mathcal{L}}_U \wedge \tilde{\mathcal{R}}_U, \quad \tilde{\mathcal{D}}_U = \tilde{\mathcal{L}}_U \vee \tilde{\mathcal{R}}_U$.
- A semigroup S with $U \subseteq E(S)$ is said to be **weakly U -abundant** if each $\tilde{\mathcal{L}}_U$ -class and each $\tilde{\mathcal{R}}_U$ -class contains an idempotent in U .
- A weakly U -abundant semigroup S satisfies **Congruence Condition (C)** if $\tilde{\mathcal{L}}_U$ is a right congruence and $\tilde{\mathcal{R}}_U$ is a left congruence.
- A weakly E -abundant semigroup (S) is an **Ehresmann semigroup** if S satisfies (C) and E is a semilattice.
- A **weakly U -concordant semigroup** is a weakly U -abundant semigroup with (C) and $\langle U \rangle$ being a regular semigroup whose set of idempotents is U .



Definition

- Let $U \subseteq E(S)$. For any $a, b \in S$,
 $a \tilde{\mathcal{L}}_U b \Leftrightarrow (\forall e \in U)(ae = a \text{ if and only if } be = b)$,
 $a \tilde{\mathcal{R}}_U b \Leftrightarrow (\forall e \in U)(ea = a \text{ if and only if } eb = b)$,
 $\tilde{\mathcal{H}}_U = \tilde{\mathcal{L}}_U \wedge \tilde{\mathcal{R}}_U, \quad \tilde{\mathcal{D}}_U = \tilde{\mathcal{L}}_U \vee \tilde{\mathcal{R}}_U$.
- A semigroup S with $U \subseteq E(S)$ is said to be **weakly U -abundant** if each $\tilde{\mathcal{L}}_U$ -class and each $\tilde{\mathcal{R}}_U$ -class contains an idempotent in U .
- A weakly U -abundant semigroup S satisfies **Congruence Condition (C)** if $\tilde{\mathcal{L}}_U$ is a right congruence and $\tilde{\mathcal{R}}_U$ is a left congruence.
- A weakly E -abundant semigroup (S) is an **Ehresmann semigroup** if S satisfies (C) and E is a semilattice.
- A **weakly U -concordant semigroup** is a weakly U -abundant semigroup with (C) and $\langle U \rangle$ being a regular semigroup whose set of idempotents is U .



Definition

- Let $U \subseteq E(S)$. For any $a, b \in S$,
 $a \tilde{\mathcal{L}}_U b \Leftrightarrow (\forall e \in U)(ae = a \text{ if and only if } be = b)$,
 $a \tilde{\mathcal{R}}_U b \Leftrightarrow (\forall e \in U)(ea = a \text{ if and only if } eb = b)$,
 $\tilde{\mathcal{H}}_U = \tilde{\mathcal{L}}_U \wedge \tilde{\mathcal{R}}_U, \quad \tilde{\mathcal{D}}_U = \tilde{\mathcal{L}}_U \vee \tilde{\mathcal{R}}_U$.
- A semigroup S with $U \subseteq E(S)$ is said to be **weakly U -abundant** if each $\tilde{\mathcal{L}}_U$ -class and each $\tilde{\mathcal{R}}_U$ -class contains an idempotent in U .
- A weakly U -abundant semigroup S satisfies **Congruence Condition (C)** if $\tilde{\mathcal{L}}_U$ is a right congruence and $\tilde{\mathcal{R}}_U$ is a left congruence.
- A weakly E -abundant semigroup (S) is an **Ehresmann semigroup** if S satisfies (C) and E is a semilattice.
- A **weakly U -concordant semigroup** is a weakly U -abundant semigroup with (C) and $\langle U \rangle$ being a regular semigroup whose set of idempotents is U .



Definition

- Let $U \subseteq E(S)$. For any $a, b \in S$,
 $a \tilde{\mathcal{L}}_U b \Leftrightarrow (\forall e \in U)(ae = a \text{ if and only if } be = b)$,
 $a \tilde{\mathcal{R}}_U b \Leftrightarrow (\forall e \in U)(ea = a \text{ if and only if } eb = b)$,
 $\tilde{\mathcal{H}}_U = \tilde{\mathcal{L}}_U \wedge \tilde{\mathcal{R}}_U, \quad \tilde{\mathcal{D}}_U = \tilde{\mathcal{L}}_U \vee \tilde{\mathcal{R}}_U$.
- A semigroup S with $U \subseteq E(S)$ is said to be **weakly U -abundant** if each $\tilde{\mathcal{L}}_U$ -class and each $\tilde{\mathcal{R}}_U$ -class contains an idempotent in U .
- A weakly U -abundant semigroup S satisfies **Congruence Condition (C)** if $\tilde{\mathcal{L}}_U$ is a right congruence and $\tilde{\mathcal{R}}_U$ is a left congruence.
- A weakly E -abundant semigroup (S) is an **Ehresmann semigroup** if S satisfies (C) and E is a semilattice.
- A **weakly U -concordant semigroup** is a weakly U -abundant semigroup with (C) and $\langle U \rangle$ being a regular semigroup whose set of idempotents is U .



Ehresmann semigroups \rightarrow Categories

Let S be an Ehresmann semigroup with distinguished semilattice of idempotents E .

$$C(S) = (S, \cdot)$$

where \cdot is defined by the rule that for any $x, y \in S$,

$$x \cdot y = \begin{cases} xy & \text{if } x^* = y^\dagger \\ \text{undefined} & \text{otherwise.} \end{cases}$$

where xy is the product of x and y in S .

Partial orders $x \leq_r y \Leftrightarrow x = x^\dagger y$; $x \leq_l y \Leftrightarrow x = yx^*$

Fact 6: The set $(C(S), \cdot, \leq_r, \leq_l)$ forms an Ehresmann category.

Ehresmann semigroups \rightarrow Categories

Let S be an Ehresmann semigroup with distinguished semilattice of idempotents E .

$$C(S) = (S, \cdot)$$

where \cdot is defined by the rule that for any $x, y \in S$,

$$x \cdot y = \begin{cases} xy & \text{if } x^* = y^\dagger \\ \text{undefined} & \text{otherwise.} \end{cases}$$

where xy is the product of x and y in S .

Partial orders $x \leq_r y \Leftrightarrow x = x^\dagger y$; $x \leq_l y \Leftrightarrow x = yx^*$

Fact 6: The set $(C(S), \cdot, \leq_r, \leq_l)$ forms an Ehresmann category.

Let S be an Ehresmann semigroup with distinguished semilattice of idempotents E .

$$C(S) = (S, \cdot)$$

where \cdot is defined by the rule that for any $x, y \in S$,

$$x \cdot y = \begin{cases} xy & \text{if } x^* = y^\dagger \\ \text{undefined} & \text{otherwise.} \end{cases}$$

where xy is the product of x and y in S .

Partial orders $x \leq_r y \Leftrightarrow x = x^\dagger y$; $x \leq_l y \Leftrightarrow x = yx^*$

Fact 6: The set $(C(S), \cdot, \leq_r, \leq_l)$ forms an Ehresmann category.

Categories \rightarrow Ehresmann semigroups

Let $(C, \cdot, \leq_r, \leq_l)$ be an Ehresmann category. We define a *pseudo-product* \otimes on C by the rule that

$$a \otimes b = (a | \mathbf{r}(a) \wedge \mathbf{d}(b)) \cdot (\mathbf{r}(a) \wedge \mathbf{d}(b) | b).$$

Fact 7: The set (C, \otimes) is an Ehresmann semigroup.

Theorem: The category of Ehresmann semigroups and admissible homomorphisms is isomorphic to the category of Ehresmann categories and strongly ordered functors. (M.V.Lawson, 1989)



Categories \rightarrow Ehresmann semigroups

Let $(C, \cdot, \leq_r, \leq_l)$ be an Ehresmann category. We define a *pseudo-product* \otimes on C by the rule that

$$a \otimes b = (a|r(a) \wedge \mathbf{d}(b)) \cdot (r(a) \wedge \mathbf{d}(b)|b).$$

Fact 7: The set (C, \otimes) is an Ehresmann semigroup.

Theorem: The category of Ehresmann semigroups and admissible homomorphisms is isomorphic to the category of Ehresmann categories and strongly ordered functors. (M.V.Lawson, 1989)



Let $(C, \cdot, \leq_r, \leq_l)$ be an Ehresmann category. We define a *pseudo-product* \otimes on C by the rule that

$$a \otimes b = (a|r(a) \wedge \mathbf{d}(b)) \cdot (r(a) \wedge \mathbf{d}(b)|b).$$

Fact 7: The set (C, \otimes) is an Ehresmann semigroup.

Theorem: The category of Ehresmann semigroups and admissible homomorphisms is isomorphic to the category of Ehresmann categories and strongly ordered functors. (M.V.Lawson, 1989)



Proper categories

Let C be a category over a regular biordered set E . Choose only one element from each \mathcal{L} -class and each \mathcal{R} -class of E containing e , denoted by e^* and e^+ , respectively. Suppose also

- 1 if $e \mathcal{R} f$ or $e \mathcal{L} f$, then $\exists [e, f]$ in $\text{Hom}(e, f)$ s.t. $[e, e] = 1_e$;
- 2 if $x \in \mathcal{C}$, $h \in E$ and $h \omega^l \mathbf{d}_x$, then there exists a **restriction** $h|_x$ such that $\mathbf{d}_{h|_x} = h$ and $\mathbf{r}_{h|_x} \omega^l \mathbf{r}_x$, in particular, $\mathbf{r}_{\mathbf{d}_x|_x} \mathcal{L} \mathbf{r}_x$ and $x \cdot [\mathbf{r}_x, \mathbf{r}_{\mathbf{d}_x|_x}] = \mathbf{d}_x|_x$;
- 3 if $e \mathcal{R} f \mathcal{R} g$ or $e \mathcal{L} f \mathcal{L} g$, then $[e, f] \cdot [f, g] = [e, g]$;
- 4 if $g \omega e$ and $e \mathcal{R} f$ or $e \mathcal{L} f$, then $g|[e, f] = [g, gf] \cdot [gf, (gf)^*]$;
- 5 if $e \omega^l f \omega^l \mathbf{d}_x$, then $e|(f|_x) = e|_x$;
- 6 if $h \omega^l \mathbf{d}_x$ and $\exists x \cdot y$, then $h|(x \cdot y) = (h|_x) \cdot (g|_y)$, where $g = \mathbf{r}_{h|_x}$.

Then C is a **proper category**.

Proper categories

Let C be a category over a regular biordered set E . Choose only one element from each \mathcal{L} -class and each \mathcal{R} -class of E containing e , denoted by e^* and e^+ , respectively. Suppose also

- 1 if $e \mathcal{R} f$ or $e \mathcal{L} f$, then $\exists [e, f]$ in $\text{Hom}(e, f)$ s.t. $[e, e] = 1_e$;
- 2 if $x \in \mathcal{C}$, $h \in E$ and $h \omega^l \mathbf{d}_x$, then there exists a **restriction** $h|_x$ such that $\mathbf{d}_{h|_x} = h$ and $\mathbf{r}_{h|_x} \omega^l \mathbf{r}_x$, in particular, $\mathbf{r}_{\mathbf{d}_x|_x} \mathcal{L} \mathbf{r}_x$ and $x \cdot [\mathbf{r}_x, \mathbf{r}_{\mathbf{d}_x|_x}] = \mathbf{d}_x|_x$;
- 3 if $e \mathcal{R} f \mathcal{R} g$ or $e \mathcal{L} f \mathcal{L} g$, then $[e, f] \cdot [f, g] = [e, g]$;
- 4 if $g \omega e$ and $e \mathcal{R} f$ or $e \mathcal{L} f$, then $g|[e, f] = [g, gf] \cdot [gf, (gf)^*]$;
- 5 if $e \omega^l f \omega^l \mathbf{d}_x$, then $e|(f|_x) = e|_x$;
- 6 if $h \omega^l \mathbf{d}_x$ and $\exists x \cdot y$, then $h|(x \cdot y) = (h|_x) \cdot (g|_y)$, where $g = \mathbf{r}_{h|_x}$.

Then C is a **proper category**.

Proper categories

Let C be a category over a regular biordered set E . Choose only one element from each \mathcal{L} -class and each \mathcal{R} -class of E containing e , denoted by e^* and e^+ , respectively. Suppose also

- 1 if $e \mathcal{R} f$ or $e \mathcal{L} f$, then $\exists [e, f]$ in $\text{Hom}(e, f)$ s.t. $[e, e] = 1_e$;
- 2 if $x \in \mathcal{C}$, $h \in E$ and $h \omega^l \mathbf{d}_x$, then there exists a **restriction** $h|_x$ such that $\mathbf{d}_{h|_x} = h$ and $\mathbf{r}_{h|_x} \omega^l \mathbf{r}_x$, in particular, $\mathbf{r}_{\mathbf{d}_x|_x} \mathcal{L} \mathbf{r}_x$ and $x \cdot [\mathbf{r}_x, \mathbf{r}_{\mathbf{d}_x|_x}] = \mathbf{d}_x|_x$;
- 3 if $e \mathcal{R} f \mathcal{R} g$ or $e \mathcal{L} f \mathcal{L} g$, then $[e, f] \cdot [f, g] = [e, g]$;
- 4 if $g \omega e$ and $e \mathcal{R} f$ or $e \mathcal{L} f$, then $g|[e, f] = [g, gf] \cdot [gf, (gf)^*]$;
- 5 if $e \omega^l f \omega^l \mathbf{d}_x$, then $e|(f|_x) = e|_x$;
- 6 if $h \omega^l \mathbf{d}_x$ and $\exists x \cdot y$, then $h|(x \cdot y) = (h|_x) \cdot (g|_y)$, where $g = \mathbf{r}_{h|_x}$.

Then C is a **proper category**.

Proper categories

Let C be a category over a regular biordered set E . Choose only one element from each \mathcal{L} -class and each \mathcal{R} -class of E containing e , denoted by e^* and e^+ , respectively. Suppose also

- 1 if $e \mathcal{R} f$ or $e \mathcal{L} f$, then $\exists [e, f]$ in $\text{Hom}(e, f)$ s.t. $[e, e] = 1_e$;
- 2 if $x \in \mathcal{C}$, $h \in E$ and $h \omega^l \mathbf{d}_x$, then there exists a **restriction** $h|_x$ such that $\mathbf{d}_{h|_x} = h$ and $\mathbf{r}_{h|_x} \omega^l \mathbf{r}_x$, in particular, $\mathbf{r}_{\mathbf{d}_x|_x} \mathcal{L} \mathbf{r}_x$ and $x \cdot [\mathbf{r}_x, \mathbf{r}_{\mathbf{d}_x|_x}] = \mathbf{d}_x|_x$;
- 3 if $e \mathcal{R} f \mathcal{R} g$ or $e \mathcal{L} f \mathcal{L} g$, then $[e, f] \cdot [f, g] = [e, g]$;
- 4 if $g \omega e$ and $e \mathcal{R} f$ or $e \mathcal{L} f$, then $g|[e, f] = [g, gf] \cdot [gf, (gf)^*]$;
- 5 if $e \omega^l f \omega^l \mathbf{d}_x$, then $e|(f|_x) = e|_x$;
- 6 if $h \omega^l \mathbf{d}_x$ and $\exists x \cdot y$, then $h|(x \cdot y) = (h|_x) \cdot (g|_y)$, where $g = \mathbf{r}_{h|_x}$.

Then C is a **proper category**.

Proper categories

Let C be a category over a regular biordered set E . Choose only one element from each \mathcal{L} -class and each \mathcal{R} -class of E containing e , denoted by e^* and e^+ , respectively. Suppose also

- 1 if $e \mathcal{R} f$ or $e \mathcal{L} f$, then $\exists [e, f]$ in $\text{Hom}(e, f)$ s.t. $[e, e] = 1_e$;
- 2 if $x \in \mathcal{C}$, $h \in E$ and $h \omega^l \mathbf{d}_x$, then there exists a **restriction** $h|_x$ such that $\mathbf{d}_{h|_x} = h$ and $\mathbf{r}_{h|_x} \omega^l \mathbf{r}_x$, in particular, $\mathbf{r}_{\mathbf{d}_x|_x} \mathcal{L} \mathbf{r}_x$ and $x \cdot [\mathbf{r}_x, \mathbf{r}_{\mathbf{d}_x|_x}] = \mathbf{d}_x|_x$;
- 3 if $e \mathcal{R} f \mathcal{R} g$ or $e \mathcal{L} f \mathcal{L} g$, then $[e, f] \cdot [f, g] = [e, g]$;
- 4 if $g \omega e$ and $e \mathcal{R} f$ or $e \mathcal{L} f$, then $g|[e, f] = [g, gf] \cdot [gf, (gf)^*]$;
- 5 if $e \omega^l f \omega^l \mathbf{d}_x$, then $e|(f|_x) = e|_x$;
- 6 if $h \omega^l \mathbf{d}_x$ and $\exists x \cdot y$, then $h|(x \cdot y) = (h|_x) \cdot (g|_y)$, where $g = \mathbf{r}_{h|_x}$.

Then C is a **proper category**.

Categories \rightarrow Weakly U -concordant semigroups

Let (P, \cdot) be a strongly proper category with $\text{Ob}(P) = U$ a regular biordered set.

Suppose that $x \in P$, $h \omega^r \mathbf{d}_x$ and $k \omega^l \mathbf{r}_x$. We define that

$$h * x = [h, h\mathbf{d}_x] \cdot h\mathbf{d}_x|x, \text{ and } x * k = x|\mathbf{r}_x k \cdot [\mathbf{r}_x k, k].$$

Define a relation ρ on P by the rule that for any $x, y \in P$,

$$x \rho y \Leftrightarrow \mathbf{d}_x \mathcal{R} \mathbf{d}_y, \mathbf{r}_x \mathcal{L} \mathbf{r}_y \text{ and } x \cdot [\mathbf{r}_x, \mathbf{r}_y] = [\mathbf{d}_x, \mathbf{d}_y] \cdot y.$$

For $x, y \in P$, $h \in S(\mathbf{r}_x, \mathbf{d}_y)$, we define that

$$\bar{x} \odot \bar{y} = \overline{(x \otimes y)_h},$$

where \bar{x} denote the ρ -class of x in P and $(x \otimes y)_h = (x * h) \cdot (h * y)$.

Fact 8: The set $\mathcal{S}(P) = (P/\rho, \odot)$ forms a weakly U -concordant semigroup.

Categories \rightarrow Weakly U -concordant semigroups

Let (P, \cdot) be a strongly proper category with $\text{Ob}(P) = U$ a regular biordered set.

Suppose that $x \in P$, $h \omega^r \mathbf{d}_x$ and $k \omega^l \mathbf{r}_x$. We define that

$$h * x = [h, h\mathbf{d}_x] \cdot h\mathbf{d}_x | x, \text{ and } x * k = x | \mathbf{r}_x k \cdot [\mathbf{r}_x k, k].$$

Define a relation ρ on P by the rule that for any $x, y \in P$,

$$x \rho y \Leftrightarrow \mathbf{d}_x \mathcal{R} \mathbf{d}_y, \mathbf{r}_x \mathcal{L} \mathbf{r}_y \text{ and } x \cdot [\mathbf{r}_x, \mathbf{r}_y] = [\mathbf{d}_x, \mathbf{d}_y] \cdot y.$$

For $x, y \in P$, $h \in S(\mathbf{r}_x, \mathbf{d}_y)$, we define that

$$\bar{x} \odot \bar{y} = \overline{(x \otimes y)_h},$$

where \bar{x} denote the ρ -class of x in P and $(x \otimes y)_h = (x * h) \cdot (h * y)$.

Fact 8: The set $\mathcal{S}(P) = (P/\rho, \odot)$ forms a weakly U -concordant semigroup.

Categories \rightarrow Weakly U -concordant semigroups

Let (P, \cdot) be a strongly proper category with $\text{Ob}(P) = U$ a regular biordered set.

Suppose that $x \in P$, $h \omega^r \mathbf{d}_x$ and $k \omega^l \mathbf{r}_x$. We define that

$$h * x = [h, h\mathbf{d}_x] \cdot h\mathbf{d}_x|_x, \text{ and } x * k = x|\mathbf{r}_x k \cdot [\mathbf{r}_x k, k].$$

Define a relation ρ on P by the rule that for any $x, y \in P$,

$$x \rho y \Leftrightarrow \mathbf{d}_x \mathcal{R} \mathbf{d}_y, \mathbf{r}_x \mathcal{L} \mathbf{r}_y \text{ and } x \cdot [\mathbf{r}_x, \mathbf{r}_y] = [\mathbf{d}_x, \mathbf{d}_y] \cdot y.$$

For $x, y \in P$, $h \in S(\mathbf{r}_x, \mathbf{d}_y)$, we define that

$$\bar{x} \odot \bar{y} = \overline{(x \otimes y)_h},$$

where \bar{x} denote the ρ -class of x in P and $(x \otimes y)_h = (x * h) \cdot (h * y)$.

Fact 8: The set $\mathcal{S}(P) = (P/\rho, \odot)$ forms a weakly U -concordant semigroup.

Categories \rightarrow Weakly U -concordant semigroups

Let (P, \cdot) be a strongly proper category with $\text{Ob}(P) = U$ a regular biordered set.

Suppose that $x \in P$, $h \omega^r \mathbf{d}_x$ and $k \omega^l \mathbf{r}_x$. We define that

$$h * x = [h, h\mathbf{d}_x] \cdot h\mathbf{d}_x|x, \text{ and } x * k = x|\mathbf{r}_x k \cdot [\mathbf{r}_x k, k].$$

Define a relation ρ on P by the rule that for any $x, y \in P$,

$$x \rho y \Leftrightarrow \mathbf{d}_x \mathcal{R} \mathbf{d}_y, \mathbf{r}_x \mathcal{L} \mathbf{r}_y \text{ and } x \cdot [\mathbf{r}_x, \mathbf{r}_y] = [\mathbf{d}_x, \mathbf{d}_y] \cdot y.$$

For $x, y \in P$, $h \in S(\mathbf{r}_x, \mathbf{d}_y)$, we define that

$$\bar{x} \odot \bar{y} = \overline{(x \otimes y)_h},$$

where \bar{x} denote the ρ -class of x in P and $(x \otimes y)_h = (x * h) \cdot (h * y)$.

Fact 8: The set $\mathcal{S}(P) = (P/\rho, \odot)$ forms a weakly U -concordant semigroup.

Categories \rightarrow Weakly U -concordant semigroups

Let (P, \cdot) be a strongly proper category with $\text{Ob}(P) = U$ a regular biordered set.

Suppose that $x \in P$, $h \omega^r \mathbf{d}_x$ and $k \omega^l \mathbf{r}_x$. We define that

$$h * x = [h, h\mathbf{d}_x] \cdot h\mathbf{d}_x | x, \text{ and } x * k = x | \mathbf{r}_x k \cdot [\mathbf{r}_x k, k].$$

Define a relation ρ on P by the rule that for any $x, y \in P$,

$$x \rho y \Leftrightarrow \mathbf{d}_x \mathcal{R} \mathbf{d}_y, \mathbf{r}_x \mathcal{L} \mathbf{r}_y \text{ and } x \cdot [\mathbf{r}_x, \mathbf{r}_y] = [\mathbf{d}_x, \mathbf{d}_y] \cdot y.$$

For $x, y \in P$, $h \in S(\mathbf{r}_x, \mathbf{d}_y)$, we define that

$$\bar{x} \odot \bar{y} = \overline{(x \otimes y)_h},$$

where \bar{x} denote the ρ -class of x in P and $(x \otimes y)_h = (x * h) \cdot (h * y)$.

Fact 8: The set $\mathcal{S}(P) = (P/\rho, \odot)$ forms a weakly U -concordant semigroup.

Weakly U -concordant semigroups \rightarrow Categories

Let S be a weakly U -concordant semigroup. Set

$$\mathbf{C}(S) = \{(e, x, f) : e \tilde{\mathcal{R}}_U x \tilde{\mathcal{L}}_U f, e, f \in U\},$$

and define a partial binary operation by the rule that

$$(e, x, f) \cdot (u, y, v) = \begin{cases} (e, xy, v) & \text{if } f = u \\ \text{undefine} & \text{otherwise,} \end{cases}$$

where xy is the product of x and y in S .

If $e \mathcal{R} f$ or $e \mathcal{L} f$, then $[e, f] = (e, ef, f)$.

Pre-orders: For any $(e, x, f) \in \mathbf{C}(S)$ and $u, v \in U$ with $u \leq_l e$ and $v \leq_r f$, we define that

$$u|(e, x, f) = (u, ux, (ux)^*) \text{ and } (e, x, f)|v = ((xv)^+, xv, v).$$

Fact 9: The set $(\mathbf{C}(S), \cdot)$ forms a strongly proper category with restrictions and co-restrictions as above.

Weakly U -concordant semigroups \rightarrow Categories

Let S be a weakly U -concordant semigroup. Set

$$\mathbf{C}(S) = \{(e, x, f) : e \tilde{\mathcal{R}}_U x \tilde{\mathcal{L}}_U f, e, f \in U\},$$

and define a partial binary operation by the rule that

$$(e, x, f) \cdot (u, y, v) = \begin{cases} (e, xy, v) & \text{if } f = u \\ \text{undefine} & \text{otherwise,} \end{cases}$$

where xy is the product of x and y in S .

If $e \mathcal{R} f$ or $e \mathcal{L} f$, then $[e, f] = (e, ef, f)$.

Pre-orders: For any $(e, x, f) \in \mathbf{C}(S)$ and $u, v \in U$ with $u \leq_l e$ and $v \leq_r f$, we define that

$$u|(e, x, f) = (u, ux, (ux)^*) \text{ and } (e, x, f)|v = ((xv)^+, xv, v).$$

Fact 9: The set $(\mathbf{C}(S), \cdot)$ forms a strongly proper category with restrictions and co-restrictions as above.

Weakly U -concordant semigroups \rightarrow Categories

Let S be a weakly U -concordant semigroup. Set

$$\mathbf{C}(S) = \{(e, x, f) : e \tilde{\mathcal{R}}_U x \tilde{\mathcal{L}}_U f, e, f \in U\},$$

and define a partial binary operation by the rule that

$$(e, x, f) \cdot (u, y, v) = \begin{cases} (e, xy, v) & \text{if } f = u \\ \text{undefine} & \text{otherwise,} \end{cases}$$

where xy is the product of x and y in S .

If $e \mathcal{R} f$ or $e \mathcal{L} f$, then $[e, f] = (e, ef, f)$.

Pre-orders: For any $(e, x, f) \in \mathbf{C}(S)$ and $u, v \in U$ with $u \leq_l e$ and $v \leq_r f$, we define that

$$u|(e, x, f) = (u, ux, (ux)^*) \text{ and } (e, x, f)|v = ((xv)^+, xv, v).$$

Fact 9: The set $(\mathbf{C}(S), \cdot)$ forms a strongly proper category with restrictions and co-restrictions as above.

Weakly U -concordant semigroups \rightarrow Categories

Let S be a weakly U -concordant semigroup. Set

$$\mathbf{C}(S) = \{(e, x, f) : e \tilde{\mathcal{R}}_U x \tilde{\mathcal{L}}_U f, e, f \in U\},$$

and define a partial binary operation by the rule that

$$(e, x, f) \cdot (u, y, v) = \begin{cases} (e, xy, v) & \text{if } f = u \\ \text{undefine} & \text{otherwise,} \end{cases}$$

where xy is the product of x and y in S .

If $e \mathcal{R} f$ or $e \mathcal{L} f$, then $[e, f] = (e, ef, f)$.

Pre-orders: For any $(e, x, f) \in \mathbf{C}(S)$ and $u, v \in U$ with $u \leq_l e$ and $v \leq_r f$, we define that

$$u|(e, x, f) = (u, ux, (ux)^*) \text{ and } (e, x, f)|v = ((xv)^+, xv, v).$$

Fact 9: The set $(\mathbf{C}(S), \cdot)$ forms a strongly proper category with restrictions and co-restrictions as above.

Weakly U -concordant semigroups \rightarrow Categories

Let S be a weakly U -concordant semigroup. Set

$$\mathbf{C}(S) = \{(e, x, f) : e \tilde{\mathcal{R}}_U x \tilde{\mathcal{L}}_U f, e, f \in U\},$$

and define a partial binary operation by the rule that

$$(e, x, f) \cdot (u, y, v) = \begin{cases} (e, xy, v) & \text{if } f = u \\ \text{undefine} & \text{otherwise,} \end{cases}$$

where xy is the product of x and y in S .

If $e \mathcal{R} f$ or $e \mathcal{L} f$, then $[e, f] = (e, ef, f)$.

Pre-orders: For any $(e, x, f) \in \mathbf{C}(S)$ and $u, v \in U$ with $u \leq_l e$ and $v \leq_r f$, we define that

$$u|(e, x, f) = (u, ux, (ux)^*) \text{ and } (e, x, f)|v = ((xv)^+, xv, v).$$

Fact 9: The set $(\mathbf{C}(S), \cdot)$ forms a strongly proper category with restrictions and co-restrictions as above.

Theorem The category of weakly U -concordant semigroups and admissible morphisms is equivalent to the category of strongly proper categories and proper functors.

Theorem: The category of inductive cancellative categories is equivalent to the category of concordant semigroups. (S. Armstrong, 1988)



Theorem The category of weakly U -concordant semigroups and admissible morphisms is equivalent to the category of strongly proper categories and proper functors.

Theorem: The category of inductive cancellative categories is equivalent to the category of concordant semigroups. (S. Armstrong, 1988)

