

Two Kinds of Congruence Networks on Regular Semigroups

Ying-Ying Feng

¹ Foshan University, Guangdong, P. R. China

² University of York, York, UK

York Semigroup, York, Apr 20, 2016

Outline

- 1 Notation & terminology
- 2 \mathcal{TK} -network on inverse semigroups
- 3 \mathcal{TV} -networks on regular semigroups
- 4 other work

Terminology

$a \in S$ is regular

$$\text{— } (\exists x \in S) axa = a$$

- regular semigroups

— all elements are regular

- inverse semigroup

— a regular semigroup whose idempotents commute

- congruence

— compatible equivalence relation, i.e.

$$(\forall s, t, s', t' \in S) [(s, t) \in \rho \text{ and } (s', t') \in \rho] \Rightarrow (ss', tt') \in \rho$$

- \mathcal{P} -type congruence

— S/ρ is a \mathcal{P} -type semigroup

- kernel-trace approach

Let ρ be a congruence on S ,

$$\text{tr } \rho = \rho|_{E(S)}, \quad \text{Ker } \rho = \{x \in S \mid (\exists e \in E(S)) x \rho e\}.$$

- inverse semigroup

1974, Scheiblich $\rho = \rho_{(\text{tr } \rho, \text{Ker } \rho)}$

1978, Petrich congruence pair

Definition

The pair (K, τ) is a *congruence pair* for S if K is a normal subsemigroup of S , τ is a normal congruence on $E(S)$, and these two satisfy:

- (i) $ae \in K, e \tau a^{-1}a \Rightarrow a \in K$ ($a \in S, e \in E(S)$);
- (ii) $k \in K \Rightarrow kk^{-1} \tau k^{-1}k$.

In such a case, define a relation $\rho_{(K, \tau)}$ on S by

$$a \rho_{(K, \tau)} b \iff a^{-1}a \tau b^{-1}b, ab^{-1} \in K.$$

Theorem

Let S be an inverse semigroup. If (K, τ) is a congruence pair for S , then $\rho_{(K, \tau)}$ is the unique congruence ρ on S for which $\text{Ker } \rho = K$ and $\text{tr } \rho = \tau$. Conversely, if ρ is a congruence on S , then $(\text{Ker } \rho, \text{tr } \rho)$ is a congruence pair for S and $\rho_{(K, \tau)} = \rho$.

Characterization of congruences

Inverse semigroup

1974, Scheiblich

$$\rho = \rho_{(\text{tr } \rho, \text{Ker } \rho)}$$

1978, Petrich

congruence pair

Regular semigroup

1979, Feigenbaum

1986, Pastijn – Petrich

Definition

A pair (K, τ) is a *congruence pair* for S if

- (i) K is a normal subset of S ,
- (ii) τ is a normal equivalence on $E(S)$,
- (iii) $K \subseteq \text{Ker}(\mathcal{L}_\tau \mathcal{L}_\tau \mathcal{L} \cap \mathcal{R}_\tau \mathcal{R}_\tau \mathcal{R})^b$,
- (iv) $\tau \subseteq \text{tr } \pi_K$.

In such a case, we define

$$\rho_{(K, \tau)} = \pi_K \cap (\mathcal{L}_\tau \mathcal{L}_\tau \mathcal{L} \cap \mathcal{R}_\tau \mathcal{R}_\tau \mathcal{R})^b.$$

Theorem

Let S be a regular semigroup. If (K, τ) is a congruence pair for S , then $\rho_{(K, \tau)}$ is the unique congruence ρ on S for which $\text{Ker } \rho = K$ and $\text{tr } \rho = \tau$. Conversely, if ρ is a congruence on S , then $(\text{Ker } \rho, \text{tr } \rho)$ is a congruence pair for S and $\rho = \rho_{(K, \tau)}$.

Definition

A triple (γ, K, δ) consisting of normal equivalences $\gamma \in \mathcal{E}(S/\mathcal{L})$ and $\delta \in \mathcal{E}(S/\mathcal{R})$ and a normal subset $K \subseteq S$, is a *congruence triple* if

- (i) $\bar{\gamma} = (\bar{\gamma} \cap \bar{\delta})^b \vee \mathcal{L}$, $\bar{\delta} = (\bar{\gamma} \cap \bar{\delta})^b \vee \mathcal{R}$;
- (ii) $K \subseteq \text{Ker } \bar{\gamma}^b$, $\bar{\gamma} \subseteq \theta_K^b \vee \mathcal{L}$;
- (iii) $K \subseteq \text{Ker } \bar{\delta}^b$, $\bar{\delta} \subseteq \theta_K^b \vee \mathcal{R}$.

If this is the case, we define

$$\rho_{(\gamma, K, \delta)} = (\bar{\gamma} \cap \theta_K \cap \bar{\delta})^b.$$

Theorem

Let S be a regular semigroup. If (γ, K, δ) is a congruence triple for S , then $\rho_{(\gamma, K, \delta)}$ is the unique congruence ρ on S such that γ is the \mathcal{L} -part of ρ , $K = \text{Ker } \rho$ and δ is the \mathcal{R} -part of ρ . Conversely, if ρ is a congruence on S , then $(\gamma, K, \delta) = ((\rho \vee \mathcal{L})/\mathcal{L}, \text{Ker } \rho, (\rho \vee \mathcal{R})/\mathcal{R})$ is a congruence triple for S and $\rho = \rho_{(\gamma, K, \delta)}$.

Congruence

- kernel-trace approach

Let ρ be a congruence on S ,

$$\text{tr } \rho = \rho|_{E(S)}, \quad \text{Ker } \rho = \{x \in S \mid (\exists e \in E(S)) x \rho e\}.$$

$$\rho = \rho_{(\text{tr } \rho, \text{Ker } \rho)}.$$

- \mathcal{T} , \mathcal{K} -relation

Let $\rho, \theta \in \mathcal{C}(S)$,

$$\rho \mathcal{T} \theta \iff \text{tr } \rho = \text{tr } \theta, \quad \rho \mathcal{K} \theta \iff \text{Ker } \rho = \text{Ker } \theta,$$

$$\rho \mathcal{U} \theta \iff \rho \cap \leq = \theta \cap \leq, \quad \rho \mathcal{V} \theta \iff \rho \mathcal{U} \theta \text{ and } \rho \mathcal{K} \theta,$$

where \leq is the natural partial order on $E(S)$.

- $\mathcal{T} \cap \mathcal{K} = \varepsilon_{\mathcal{C}(S)} = \mathcal{T} \cap \mathcal{V}$

Definition

A triple (γ, π, δ) consisting of normal equivalences $\gamma \in \mathcal{E}(S/\mathcal{L})$ and $\delta \in \mathcal{E}(S/\mathcal{R})$ and a \mathcal{V} -normal congruence π on S , is a \mathcal{VT} -congruence triple if

- (i) $\bar{\gamma} = (\bar{\gamma} \cap \bar{\delta})^b \vee \mathcal{L}$, $\bar{\delta} = (\bar{\gamma} \cap \bar{\delta})^b \vee \mathcal{R}$;
- (ii) $\pi \subseteq (\bar{\gamma}^b)^\vee$, $\bar{\gamma} \subseteq \pi \vee \mathcal{L}$;
- (iii) $\pi \subseteq (\bar{\delta}^b)^\vee$, $\bar{\delta} \subseteq \pi \vee \mathcal{R}$.

If this is the case, we define

$$\rho_{(\gamma, \pi, \delta)} = (\bar{\gamma} \cap \pi \cap \bar{\delta})^b.$$

Theorem

Let S be a regular semigroup. If (γ, π, δ) is a \mathcal{VT} -congruence triple for S , then $\rho_{(\gamma, \pi, \delta)}$ is the unique congruence ρ on S such that γ is the \mathcal{L} -part of ρ , π is the \mathcal{V} -part of ρ and δ is the \mathcal{R} -part of ρ . Conversely, if ρ is a congruence on S , then $(\gamma, \pi, \delta) = ((\rho \vee \mathcal{L})/\mathcal{L}, \overline{\mathcal{V}_{S/\rho}}^b, (\rho \vee \mathcal{R})/\mathcal{R})$ is a congruence triple for S and $\rho = \rho_{(\gamma, \pi, \delta)}$.

- $\rho \mathcal{T} \theta \iff \text{tr } \rho = \text{tr } \theta, \quad \rho \mathcal{K} \theta \iff \text{Ker } \rho = \text{Ker } \theta,$
 $\rho \mathcal{U} \theta \iff \rho \cap \leq = \theta \cap \leq, \quad \mathcal{V} = \mathcal{U} \cap \mathcal{K}.$

Result

For any $\rho \in \mathcal{C}(S)$, $\rho \mathcal{T} = [\rho t, \rho T]$, $\rho \mathcal{K} = [\rho k, \rho K]$, $\rho \mathcal{U} = [\rho u, \rho U]$,
 $\rho \mathcal{V} = [\rho v, \rho V]$, where

$$\rho t = (\text{tr } \rho)^\sharp, \quad \rho T = \overline{\mathcal{H}_{S/\rho}}^b,$$

$$\rho k = \{(x, x^2) \in S \times S \mid x \in \text{Ker } \rho\}^\sharp, \quad \rho K = \theta_{\text{Ker } \rho}^b,$$


$$\rho u = (\rho \cap \leq)^\sharp, \quad \rho U = \overline{\mathcal{U}_{S/\rho}}^b,$$

$$\rho v = \rho U \vee \rho K, \quad \rho V = \rho U \cap \rho K = \overline{\mathcal{V}_{S/\rho}}^b.$$


- kernel-trace approach
- \mathcal{T} , \mathcal{K} -relation
- congruence networks
 - single out various classes of semigroups of particular interest
 - structure

Congruence network

$$\mathcal{T} \cap \mathcal{K} = \varepsilon$$




\mathcal{TK} -network of ρ



\mathcal{TK} -min network of ρ


Congruence network

$$\mathcal{T} \cap \mathcal{K} = \varepsilon$$




\mathcal{TK} -network of ρ

$$\mathcal{T} \cap \mathcal{V} = \varepsilon$$




\mathcal{TV} -network of ρ

Inverse semigroup




TK min-network of ω
 [1982, Petrich – Reilly]

Regular semigroup



TK min-network of ω
 [1988, Alimpić – Krgović]

\mathcal{TK} -network on inverse semigroup



\mathcal{TK} min-network of ω

$E\omega$ -Clifford semigroup and $E\omega$ - E -reflexive semigroup

Proposition

The following conditions on an inverse semigroup S are equivalent.

- (1) S is an $E\omega$ -Clifford semigroup;
- (2) $\sigma \cap \mathcal{L}$ is a congruence;
- (3) $\sigma \cap \mathcal{R}$ is a congruence;
- (4) $\sigma \cap \mathcal{L} = \sigma \cap \mathcal{R}$;
- (5) $\sigma \cap \mathcal{L} = \sigma \cap \mu$;
- (6) there exists an idempotent separating E -unitary congruence on S ;
- (7) $\pi \subseteq \mu$;
- (8) $\pi t = \varepsilon$;
- (9) $e\sigma$ is a Clifford semigroup for every $e \in E(S)$;
- (10) S satisfies the implication $xy = x \Rightarrow y \in E(S) \zeta$;
- (11) $E(S)\omega \subseteq E(S) \zeta$;
- (12) $\pi \cap \mathcal{F} = \varepsilon$.

Theorem

The following conditions on an inverse semigroup S are equivalent.

- (1) S is $E\omega$ - E -reflexive;
- (2) $\pi \cap \mathcal{F}$ is a congruence;
- (3) $\pi \cap \mathcal{C}$ is a congruence;
- (4) $\pi \cap \mathcal{F} = \pi \cap \tau$;
- (5) $\pi \cap \mathcal{C} = \pi \cap \tau$;
- (6) there exists an idempotent pure $E\omega$ -Clifford congruence on S ;
- (7) $\zeta \subseteq \tau$;
- (8) $\zeta k = \varepsilon$;
- (9) $e\pi$ is E -unitary for every $e \in E(S)$;
- (10) S satisfies the implication $xy = x, x\pi y \Rightarrow y \in E(S)$;
- (11) $\zeta \cap \mathcal{L} = \varepsilon$.

$E\omega$ -Clifford congruence and $E\omega$ - E -reflexive congruence

Proposition

The following statements concerning a congruence ρ on an inverse semigroup S are equivalent.

- (1) ρ is an $E\omega$ -Clifford congruence;*
- (2) $\pi_\rho \subseteq \rho T$, where π_ρ is the least E -unitary congruence on S containing ρ ;*
- (3) $\text{tr } \pi_\rho = \text{tr } \rho$.*

Proposition

The following statements concerning a congruence ρ on an inverse semigroup S are equivalent.

- (1) ρ is $E\omega$ - E -reflexive;*
- (2) $\zeta_\rho \subseteq \rho K$, where ζ_ρ is the least $E\omega$ -Clifford congruence on S containing ρ ;*
- (3) $\text{Ker } \zeta_\rho = \text{Ker } \rho$.*

Coincidences

Petrich, *Inverse Semigroups*, Table III.8.10


	ω	σ	η	ν	π	λ	ζ	χ	μ	τ
σ	$E\omega = S$									
η	no c. pr. ideals	$E\omega = S$, no c. pr. ideals								
ν	$\sigma = \eta = \omega$	no c. pr. ideals	$E_A\omega = A$ ($\forall \eta$ -cl. A)							
π	$\sigma = \eta = \omega$	$\text{tr } \pi = \omega$	$E\omega = S$							
λ	$\sigma = \eta = \omega$	$\text{tr } \pi = \omega$	$E_A\omega = A$ ($\forall \eta$ -cl. A)							
ζ	$\sigma = \eta = \omega$	$\text{tr } \pi = \omega$	$E_A\omega = A$ ($\forall \eta$ -cl. A)							
χ	$\sigma = \eta = \omega$	$\text{tr } \pi = \omega$	$E_A\omega = A$ ($\forall \eta$ -cl. A)							
μ	group	trivial	Clifford	semil.	$E\omega = E\zeta$ and $\text{tr } \pi = \varepsilon$					
τ	semil.	E -un.	trivial	E -refl. $\text{tr } \pi = \text{tr } \eta$	E -un. E -disj.	E -refl. $\text{tr } \tau = \text{tr } \lambda$	$E\omega$ - E -refl. $\text{tr } \tau = \text{tr } \pi$	E -disj. $E\omega$ - E -refl.	E -disj. antig.	
ε	trivial	group	semil.	Clifford	E -un.	E -refl.	$E\omega$ -Clifford	$E\omega$ - E -refl.	antig.	E -disj.

● Fill


● Refine

● Add

Question



\mathcal{TK} -network on inverse semigroup



\mathcal{TK} min-network of ω


\mathcal{TV} -network of ω

$$\rho \mathcal{K} \theta \iff \text{Ker } \rho = \text{Ker } \theta$$

$$\rho \mathcal{U} \theta \iff \rho \cap \leq = \theta \cap \leq$$

$$\mathcal{V} = \mathcal{U} \cap \mathcal{K}$$


Inverse semigroup $\mathcal{V} = \varepsilon$



\mathcal{TV} -network of ω

\mathcal{TV} -network of ω


Regular semigroup




δt
completely simple
— rectangular band of groups
[1995, Wang, Q-inverse transversal]

σv
rectangular group
[1995, Wang]

Inverse semigroup




Regular semigroup




\mathcal{TV} -network of ω

Regular semigroup




\mathcal{TV} -min network of ω

\mathcal{TV} -network of ω



\mathcal{TV} -min network of ω



\mathcal{TV} -network of ω

Theorem

For a congruence ρ on a regular semigroup S .

- (1) ρt is over bands $\iff \rho t = \rho \cap \tau \implies \rho$ is over E -unitary semigroups;
- (2) ρt is over rectangular bands $\iff \rho t = \rho \cap \varepsilon V \implies \rho$ is over rectangular groups;
- (3) ρv is over groups $\iff \rho v = \rho \cap \mu \implies \rho$ is over completely simple semigroups;
- (4) ρk is over groups $\iff \rho k = \rho \cap \mu \implies \rho$ is over cryptogroups.


Corollary

On a regular semigroup S , the following statements hold.


- (1) τT is over E -unitary semigroups;
- (2) $(\varepsilon V)T$ is over rectangular groups;
- (3) μV is over completely simple semigroups;
- (4) μK is over cryptogroups.

\mathcal{TV} -network of ε

Cryptogroup




\mathcal{TV} -max network of ε




\mathcal{TV} -network of ε

\mathcal{TV} -network of η





\mathcal{TV} -min network of η



- orthogroup
orthodox completely regular semigroup
- rectangular group
orthodox completely simple semigroup;
equivalently, a direct product of a rectangular band and a group

S	orthodox	orthogroup	rectangular group	band
\iff	$\varepsilon V = \gamma$	$\tau V = \nu$	$\varepsilon V = \sigma$	$\varepsilon V = \eta$
$\iff \forall \rho \in \mathcal{C}(S)$	$\rho V = \rho \vee \gamma$	$\rho V = \rho \vee \nu$	$\rho V = \rho \vee \sigma$	$\rho V = \rho \vee \eta$
$\iff \rho V$ is	inverse	Clifford	group	semilattice
$\iff S$ is coextension of	inverse semigroup by rectangular bands	Clifford semigroup by rectangular bands	group by rectangular bands	



S	E -solid	CCS	CGS	completely regular
$\iff \mathcal{U}^0$	inverse	Clifford	group	semilattice
$\iff \rho U$	inverse	Clifford	group	semilattice

- E -solid

$$\mathcal{R}|_E \circ \mathcal{L}|_E = \mathcal{L}|_E \circ \mathcal{R}|_E$$

- CCS

coextensions of Clifford semigroups by completely simple semigroups

- CGS

coextensions of groups by completely simple semigroups

Congruence

- kernel-trace approach
- \mathcal{T} , \mathcal{K} -relation
- congruence networks
- operator semigroup

Operator semigroup

Four operators:

$$T : \lambda \mapsto \lambda T, \quad t : \lambda \mapsto \lambda t, \quad K : \lambda \mapsto \lambda K, \quad k : \lambda \mapsto \lambda k.$$


$$\Gamma = \{T, t, K, k\}$$

- \mathcal{TK} -network

$$\rho, \rho T, \rho t, \rho K, \rho k, \rho TK, \rho Tk, \dots$$

- Γ^+ , \mathcal{TK} -operator semigroup [1992, Petrich]
 - relation Σ — valid in all networks of congruences
 - $\Gamma^+ / \Sigma^\#$

\mathcal{TK} -operator semigroup for Clifford semigroups



\mathcal{TK} -network of ω

Lemma

Operators Γ satisfy the following relations Σ .

- (1) $K^2 = kK = K, \quad k^2 = Kk = k,$
 $t^2 = Tt = t, \quad T^2 = tT = T;$
- (2) $KTK = TKT = TK, \quad tkt = ktk = kt;$
- (3) $tKt = tK;$
- (4) $kT = Tk.$

Denote

$$\begin{aligned} \varepsilon &= kt, & \tau &= ktK, & \tau \vee \eta &= ktKT, & \eta &= kT, \\ \omega &= TK, & \sigma &= TKt, & \sigma \cap \eta &= TKtk. \end{aligned}$$

Let

$$\Delta = \{\varepsilon, \sigma, \eta, \tau, \sigma \cap \eta, \tau \vee \eta, \omega\}.$$

Theorem

Let S be a Clifford semigroup. The set

$$\Omega = \{ K, \quad KT, \quad Kt, \quad KtK, \quad Ktk, \quad KtKT, \quad k, \\ t, \quad tk, \quad tK, \quad tKT, \quad T \} \cup \Delta$$

is a system of representatives for the congruence on Γ^+ generated by the relations Σ .

Theorem

The \mathcal{TK} -operator semigroup for Clifford semigroups is $\Gamma^+ / \Sigma^\#$.

other \mathcal{TK} -operator semigroups

- completely simple semigroup [1994, Petrich]
- cryptogroup [2000, Wang]
- bisimple ω -semigroup [2000, Wang]
- E -unitary completely regular semigroup [2001, Luo – Wang]
- free monogenic inverse semigroup [2014, Long – Wang]

- congruence

$$\rho = \rho(\text{tr } \rho, \text{Ker } \rho)$$

- $\mathcal{T}, \mathcal{K}, \mathcal{U}, \mathcal{V}$

$$\rho \mathcal{T} \theta \iff \text{tr } \rho = \text{tr } \theta,$$

$$\rho \mathcal{K} \theta \iff \text{Ker } \rho = \text{Ker } \theta,$$


$$\rho \mathcal{U} \theta \iff \rho \cap \leq = \theta \cap \leq,$$

$$\mathcal{V} = \mathcal{U} \cap \mathcal{K}.$$

- congruence network

- operator semigroup

$$\Gamma^+ / \Sigma^\sharp, \text{ where } \Gamma = \{T, t, K, k\}.$$



\mathcal{TK} -min network of ω